

2014 KENYA DHS
Demographic and Health Survey

MOMBASA


	Mombasa	Kenya total
Fertility and Family Planning		
Total fertility rate (number of children per woman)	3.2	3.9
Current use of *any* method of family planning (% of currently married women age 15-49)	55	58
Current use of any modern method of family planning (% of currently married women age 15-49)	44	53
Maternal and Child Health		
Maternity care		
Pregnant women who received antenatal care from a skilled provider[1] (%)	99	96
Pregnant women who received 4+ ANC visits (%)	65	58
Births assisted by a skilled provider[1] (%)	83	62
Births delivered in a health facility (%)	82	61
Child vaccination		
Children 12–23 months fully vaccinated[2] (%)	68	68
Nutrition		
Children under 5 years who are stunted (moderate or severe) (%)	21	26
Children under 5 years who are wasted (moderate or severe) (%)	4	4
Children under 5 years who are underweight (%)	10	11
Children under 5 years who are overweight or obese (%)	4	4
Malaria Prevention and Treatment		
Households with at least one insecticide-treated net[3] (ITN) (%)	57	59
Households with at least one ITN for every 2 persons in the household (universal coverage) (%)	39	34
Children under age 5 who slept under an ITN the night before the survey (%)	62	54
Pregnant women who slept under an ITN the night before the survey (%)	(55)	51
Pregnant women who received 3+ doses of SP/Fansidar to prevent malaria during pregnancy, at least one dose was received during an ANC visit (IPTp) (%)	25	10
Children under age 5 with fever in the 2 weeks before the survey	23	24
Among children under age 5 with fever, percentage who took any ACT	14	23
HIV/AIDS-related Knowledge		
Women 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	83	77
Men 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	96	85
Women 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	58	53
Men 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	40	46

Note: Figures in parentheses are based on 25-49 unweighted cases; an asterisk indicates that data are based on fewer than 25 unweighted cases and have been suppressed. [1]Skilled provider includes doctor, nurse, or midwife. [2]Fully vaccinated includes BCG, measles, three doses each of DPT and polio vaccine (excluding polio vaccine given at birth) and pneumococcal. [3]An insecticide-treated net (ITN) is (1) a factory-treated net that does not require any further treatment, or (2) a net that has been soaked with insecticide within the past 6 months.

Coast: Key Indicators from the 2014 KDHS


Total Fertility Rate

Average number of live births per woman age 15-49




Use of Modern Methods of Family Planning

Percent of currently married women using a modern method of contraception




Antenatal Care: 4+ visits

Among women with a live birth in the 5 years before the survey, percent with 4+ ANC visits for their last live birth




Delivery in a Health Facility

Percentage of live births delivered in a health facility




Full Vaccination Rate

Percent of children age 12-23 months who have received all basic vaccinations plus the pneumococcal vaccine



ITN Access

Percentage of households with at least one insecticide-treated net (ITN) for every two persons who stayed in the household last night




The 2014 Kenya Demographic and Health Survey (2014 KDHS) was implemented by the Kenya National Bureau of Statistics from May 2014 to October 2014 in partnership with the Ministry of Health, the National AIDS Control Council (NACC), the National Council for Population and Development (NCPD), and the Kenya Medical Research Institute (KEMRI). Funding for the KDHS was provided by the Government of Kenya with support from the United States Agency for International Development (USAID), the United Nations Population Fund (UNFPA), the United Kingdom Department for International Development (DfID), the World Bank, the Danish International Development Agency (DANIDA), the United Nations Children's Fund (UNICEF), the German Development Bank (KfW), the Clinton Health Access Initiative (CHAI), the World Food Programme (WFP), and the Micronutrient Initiative (MI). ICF International provided technical assistance as well as funding to the project through The DHS Program, a USAID-funded project providing support and technical assistance in the implementation of population and health surveys in countries worldwide.

Republic of Kenya

Additional information about the 2014 KDHS may be obtained from the Kenya National Bureau of Statistics (KNBS), P.O. Box 30266-00100 GPO Nairobi, Kenya; telephone (Nairobi): 3317586/8, 3317612/22, 3317623, 3317651; fax: 3315977; e-mail: directorgeneral@knbs.or.ke, info@knbs.or.ke; website: www.knbs.or.ke.

KNBS KENYA NATIONAL BUREAU OF STATISTICS *Keeping you informed*

Information on The DHS Program may be obtained from ICF International, 530 Gaither Road, Suite 500, Rockville MD, 20850, USA; telephone: 301-407-6500; fax: 301.407.6501; e-mail: info@DHSprogram.com; website: www.DHSprogram.com.



KWALE

2014 KENYA DHS
Demographic and Health Survey


	Kwale	Kenya total
Fertility and Family Planning		
Total fertility rate (number of children per woman)	4.7	3.9
Current use of *any* method of family planning (% of currently married women age 15-49)	42	58
Current use of any modern method of family planning (% of currently married women age 15-49)	38	53
Maternal and Child Health		
Maternity care		
Pregnant women who received antenatal care from a skilled provider[1] (%)	96	96
Pregnant women who received 4+ ANC visits (%)	60	58
Births assisted by a skilled provider[1] (%)	50	62
Births delivered in a health facility (%)	49	61
Child vaccination		
Children 12–23 months fully vaccinated[2] (%)	82	68
Nutrition		
Children under 5 years who are stunted (moderate or severe) (%)	30	26
Children under 5 years who are wasted (moderate or severe) (%)	4	4
Children under 5 years who are underweight (%)	12	11
Children under 5 years who are overweight or obese (%)	4	4
Malaria Prevention and Treatment		
Households with at least one insecticide-treated net[3] (ITN) (%)	81	59
Households with at least one ITN for every 2 persons in the household (universal coverage) (%)	48	34
Children under age 5 who slept under an ITN the night before the survey (%)	72	54
Pregnant women who slept under an ITN the night before the survey (%)	(85)	51
Pregnant women who received 3+ doses of SP/Fansidar to prevent malaria during pregnancy, at least one dose was received during an ANC visit (IPTp) (%)	52	10
Children under age 5 with fever in the 2 weeks before the survey	26	24
Among children under age 5 with fever, percentage who took any ACT	30	23
HIV/AIDS-related Knowledge		
Women 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	56	77
Men 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	67	85
Women 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	52	53
Men 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	37	46

Note: Figures in parentheses are based on 25-49 unweighted cases; an asterisk indicates that data are based on fewer than 25 unweighted cases and have been suppressed. [1]Skilled provider includes doctor, nurse, or midwife. [2]Fully vaccinated includes BCG, measles, three doses each of DPT and polio vaccine (excluding polio vaccine given at birth) and pneumococcal. [3]An insecticide-treated net (ITN) is (1) a factory-treated net that does not require any further treatment, or (2) a net that has been soaked with insecticide within the past 6 months.

Coast: Key Indicators from the 2014 KDHS


Total Fertility Rate

Average number of live births per woman age 15-49




Use of Modern Methods of Family Planning

Percent of currently married women using a modern method of contraception




Antenatal Care: 4+ visits

Among women with a live birth in the 5 years before the survey, percent with 4+ ANC visits for their last live birth




Delivery in a Health Facility

Percentage of live births delivered in a health facility




Full Vaccination Rate

Percent of children age 12-23 months who have received all basic vaccinations plus the pneumococcal vaccine



ITN Access

Percentage of households with at least one insecticide-treated net (ITN) for every two persons who stayed in the household last night




The 2014 Kenya Demographic and Health Survey (2014 KDHS) was implemented by the Kenya National Bureau of Statistics from May 2014 to October 2014 in partnership with the Ministry of Health, the National AIDS Control Council (NACC), the National Council for Population and Development (NCPD), and the Kenya Medical Research Institute (KEMRI). Funding for the KDHS was provided by the Government of Kenya with support from the United States Agency for International Development (USAID), the United Nations Population Fund (UNFPA), the United Kingdom Department for International Development (DfID), the World Bank, the Danish International Development Agency (DANIDA), the United Nations Children's Fund (UNICEF), the German Development Bank (KfW), the Clinton Health Access Initiative (CHAI), the World Food Programme (WFP), and the Micronutrient Initiative (MI). ICF International provided technical assistance as well as funding to the project through The DHS Program, a USAID-funded project providing support and technical assistance in the implementation of population and health surveys in countries worldwide.

Republic of Kenya

Additional information about the 2014 KDHS may be obtained from the Kenya National Bureau of Statistics (KNBS), P.O. Box 30266-00100 GPO Nairobi, Kenya; telephone (Nairobi): 3317586/8, 3317612/22, 3317623, 3317651; fax: 3315977; e-mail: directorgeneral@knbs.or.ke, info@knbs.or.ke; website: www.knbs.or.ke.

KNBS KENYA NATIONAL BUREAU OF STATISTICS *Keeping you informed*

Information on The DHS Program may be obtained from ICF International, 530 Gaither Road, Suite 500, Rockville MD, 20850, USA; telephone: 301-407-6500; fax: 301.407.6501; e-mail: info@DHSprogram.com; website: www.DHSprogram.com.



KILIFI

2014 KENYA DHS
Demographic and Health Survey


	Kilifi	Kenya total
Fertility and Family Planning		
Total fertility rate (number of children per woman)	5.1	3.9
Current use of *any* method of family planning (% of currently married women age 15-49)	34	58
Current use of any modern method of family planning (% of currently married women age 15-49)	33	53
Maternal and Child Health		
Maternity care		
Pregnant women who received antenatal care from a skilled provider[1] (%)	98	96
Pregnant women who received 4+ ANC visits (%)	65	58
Births assisted by a skilled provider[1] (%)	52	62
Births delivered in a health facility (%)	53	61
Child vaccination		
Children 12–23 months fully vaccinated[2] (%)	72	68
Nutrition		
Children under 5 years who are stunted (moderate or severe) (%)	40	26
Children under 5 years who are wasted (moderate or severe) (%)	4	4
Children under 5 years who are underweight (%)	17	11
Children under 5 years who are overweight or obese (%)	3	4
Malaria Prevention and Treatment		
Households with at least one insecticide-treated net[3] (ITN) (%)	73	59
Households with at least one ITN for every 2 persons in the household (universal coverage) (%)	42	34
Children under age 5 who slept under an ITN the night before the survey (%)	64	54
Pregnant women who slept under an ITN the night before the survey (%)	58	51
Pregnant women who received 3+ doses of SP/Fansidar to prevent malaria during pregnancy, at least one dose was received during an ANC visit (IPTp) (%)	29	10
Children under age 5 with fever in the 2 weeks before the survey	31	24
Among children under age 5 with fever, percentage who took any ACT	1	23
HIV/AIDS-related Knowledge		
Women 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	73	77
Men 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	44	85
Women 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	54	53
Men 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	42	46

Note: Figures in parentheses are based on 25-49 unweighted cases; an asterisk indicates that data are based on fewer than 25 unweighted cases and have been suppressed. [1]Skilled provider includes doctor, nurse, or midwife. [2]Fully vaccinated includes BCG, measles, three doses each of DPT and polio vaccine (excluding polio vaccine given at birth) and pneumococcal. [3]An insecticide-treated net (ITN) is (1) a factory-treated net that does not require any further treatment, or (2) a net that has been soaked with insecticide within the past 6 months.

Coast: Key Indicators from the 2014 KDHS


Total Fertility Rate

Average number of live births per woman age 15-49




Use of Modern Methods of Family Planning

Percent of currently married women using a modern method of contraception




Antenatal Care: 4+ visits

Among women with a live birth in the 5 years before the survey, percent with 4+ ANC visits for their last live birth




Delivery in a Health Facility

Percentage of live births delivered in a health facility




Full Vaccination Rate

Percent of children age 12-23 months who have received all basic vaccinations plus the pneumococcal vaccine



ITN Access

Percentage of households with at least one insecticide-treated net (ITN) for every two persons who stayed in the household last night




The 2014 Kenya Demographic and Health Survey (2014 KDHS) was implemented by the Kenya National Bureau of Statistics from May 2014 to October 2014 in partnership with the Ministry of Health, the National AIDS Control Council (NACC), the National Council for Population and Development (NCPD), and the Kenya Medical Research Institute (KEMRI). Funding for the KDHS was provided by the Government of Kenya with support from the United States Agency for International Development (USAID), the United Nations Population Fund (UNFPA), the United Kingdom Department for International Development (DfID), the World Bank, the Danish International Development Agency (DANIDA), the United Nations Children's Fund (UNICEF), the German Development Bank (KfW), the Clinton Health Access Initiative (CHAI), the World Food Programme (WFP), and the Micronutrient Initiative (MI). ICF International provided technical assistance as well as funding to the project through The DHS Program, a USAID-funded project providing support and technical assistance in the implementation of population and health surveys in countries worldwide.

Republic of Kenya

Additional information about the 2014 KDHS may be obtained from the Kenya National Bureau of Statistics (KNBS), P.O. Box 30266-00100 GPO Nairobi, Kenya; telephone (Nairobi): 3317586/8, 3317612/22, 3317623, 3317651; fax: 3315977; e-mail: directorgeneral@knbs.or.ke, info@knbs.or.ke; website: www.knbs.or.ke.

KNBS KENYA NATIONAL BUREAU OF STATISTICS *Keeping you informed*

Information on The DHS Program may be obtained from ICF International, 530 Gaither Road, Suite 500, Rockville MD, 20850, USA; telephone: 301-407-6500; fax: 301.407.6501; e-mail: info@DHSprogram.com; website: www.DHSprogram.com.



2014 Kenya DHS
Demographic and Health Survey

Tana River


	Tana River	Kenya total
Fertility and Family Planning		
Total fertility rate (number of children per woman)	5.8	3.9
Current use of *any* method of family planning (% of currently married women age 15-49)	29	58
Current use of any modern method of family planning (% of currently married women age 15-49)	21	53
Maternal and Child Health		
Maternity care		
Pregnant women who received antenatal care from a skilled provider[1] (%)	94	96
Pregnant women who received 4+ ANC visits (%)	51	58
Births assisted by a skilled provider[1] (%)	32	62
Births delivered in a health facility (%)	32	61
Child vaccination		
Children 12–23 months fully vaccinated[2] (%)	61	68
Nutrition		
Children under 5 years who are stunted (moderate or severe) (%)	28	26
Children under 5 years who are wasted (moderate or severe) (%)	6	4
Children under 5 years who are underweight (%)	19	11
Children under 5 years who are overweight or obese (%)	1	4
Malaria Prevention and Treatment		
Households with at least one insecticide-treated net[3] (ITN) (%)	65	59
Households with at least one ITN for every 2 persons in the household (universal coverage) (%)	33	34
Children under age 5 who slept under an ITN the night before the survey (%)	57	54
Pregnant women who slept under an ITN the night before the survey (%)	48	51
Pregnant women who received 3+ doses of SP/Fansidar to prevent malaria during pregnancy, at least one dose was received during an ANC visit (IPTp) (%)	21	10
Children under age 5 with fever in the 2 weeks before the survey	27	24
Among children under age 5 with fever, percentage who took any ACT	10	23
HIV/AIDS-related Knowledge		
Women 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	58	77
Men 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	71	85
Women 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	42	53
Men 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	36	46

Note: Figures in parentheses are based on 25-49 unweighted cases; an asterisk indicates that data are based on fewer than 25 unweighted cases and have been suppressed. [1]Skilled provider includes doctor, nurse, or midwife. [2]Fully vaccinated includes BCG, measles, three doses each of DPT and polio vaccine (excluding polio vaccine given at birth) and pneumococcal. [3]An insecticide-treated net (ITN) is (1) a factory-treated net that does not require any further treatment, or (2) a net that has been soaked with insecticide within the past 6 months.

Coast: Key Indicators from the 2014 KDHS


Total Fertility Rate

Average number of live births per woman age 15-49




Use of Modern Methods of Family Planning

Percent of currently married women using a modern method of contraception




Antenatal Care: 4+ visits

Among women with a live birth in the 5 years before the survey, percent with 4+ ANC visits for their last live birth




Delivery in a Health Facility

Percentage of live births delivered in a health facility




Full Vaccination Rate

Percent of children age 12-23 months who have received all basic vaccinations plus the pneumococcal vaccine



ITN Access

Percentage of households with at least one insecticide-treated net (ITN) for every two persons who stayed in the household last night




The 2014 Kenya Demographic and Health Survey (2014 KDHS) was implemented by the Kenya National Bureau of Statistics from May 2014 to October 2014 in partnership with the Ministry of Health, the National AIDS Control Council (NACC), the National Council for Population and Development (NCPD), and the Kenya Medical Research Institute (KEMRI). Funding for the KDHS was provided by the Government of Kenya with support from the United States Agency for International Development (USAID), the United Nations Population Fund (UNFPA), the United Kingdom Department for International Development (DfID), the World Bank, the Danish International Development Agency (DANIDA), the United Nations Children's Fund (UNICEF), the German Development Bank (KfW), the Clinton Health Access Initiative (CHAI), the World Food Programme (WFP), and the Micronutrient Initiative (MI). ICF International provided technical assistance as well as funding to the project through The DHS Program, a USAID-funded project providing support and technical assistance in the implementation of population and health surveys in countries worldwide.

Republic of Kenya

Additional information about the 2014 KDHS may be obtained from the Kenya National Bureau of Statistics (KNBS), P.O. Box 30266-00100 GPO Nairobi, Kenya; telephone (Nairobi): 3317586/8, 3317612/22, 3317623, 3317651; fax: 3315977; e-mail: directorgeneral@knbs.or.ke, info@knbs.or.ke; website: www.knbs.or.ke.

KNBS KENYA NATIONAL BUREAU OF STATISTICS *Keeping you informed*

Information on The DHS Program may be obtained from ICF International, 530 Gaither Road, Suite 500, Rockville MD, 20850, USA; telephone: 301-407-6500; fax: 301.407.6501; e-mail: info@DHSprogram.com; website: www.DHSprogram.com.



2014 Kenya DHS
Demographic and Health Survey

Lamu


	Lamu	Kenya total
Fertility and Family Planning		
Total fertility rate (number of children per woman)	4.3	3.9
Current use of *any* method of family planning (% of currently married women age 15-49)	42	58
Current use of any modern method of family planning (% of currently married women age 15-49)	40	53
Maternal and Child Health		
Maternity care		
Pregnant women who received antenatal care from a skilled provider[1] (%)	96	96
Pregnant women who received 4+ ANC visits (%)	62	58
Births assisted by a skilled provider[1] (%)	47	62
Births delivered in a health facility (%)	44	61
Child vaccination		
Children 12–23 months fully vaccinated[2] (%)	64	68
Nutrition		
Children under 5 years who are stunted (moderate or severe) (%)	29	26
Children under 5 years who are wasted (moderate or severe) (%)	4	4
Children under 5 years who are underweight (%)	12	11
Children under 5 years who are overweight or obese (%)	2	4
Malaria Prevention and Treatment		
Households with at least one insecticide-treated net[3] (ITN) (%)	71	59
Households with at least one ITN for every 2 persons in the household (universal coverage) (%)	51	34
Children under age 5 who slept under an ITN the night before the survey (%)	58	54
Pregnant women who slept under an ITN the night before the survey (%)	(64)	51
Pregnant women who received 3+ doses of SP/Fansidar to prevent malaria during pregnancy, at least one dose was received during an ANC visit (IPTp) (%)	36	10
Children under age 5 with fever in the 2 weeks before the survey	22	24
Among children under age 5 with fever, percentage who took any ACT	1	23
HIV/AIDS-related Knowledge		
Women 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	68	77
Men 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	84	85
Women 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	45	53
Men 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	41	46

Note: Figures in parentheses are based on 25-49 unweighted cases; an asterisk indicates that data are based on fewer than 25 unweighted cases and have been suppressed. [1]Skilled provider includes doctor, nurse, or midwife. [2]Fully vaccinated includes BCG, measles, three doses each of DPT and polio vaccine (excluding polio vaccine given at birth) and pneumococcal. [3]An insecticide-treated net (ITN) is (1) a factory-treated net that does not require any further treatment, or (2) a net that has been soaked with insecticide within the past 6 months.

Coast: Key Indicators from the 2014 KDHS


Total Fertility Rate

Average number of live births per woman age 15-49




Use of Modern Methods of Family Planning

Percent of currently married women using a modern method of contraception




Antenatal Care: 4+ visits

Among women with a live birth in the 5 years before the survey, percent with 4+ ANC visits for their last live birth




Delivery in a Health Facility

Percentage of live births delivered in a health facility




Full Vaccination Rate

Percent of children age 12-23 months who have received all basic vaccinations plus the pneumococcal vaccine



ITN Access

Percentage of households with at least one insecticide-treated net (ITN) for every two persons who stayed in the household last night




The 2014 Kenya Demographic and Health Survey (2014 KDHS) was implemented by the Kenya National Bureau of Statistics from May 2014 to October 2014 in partnership with the Ministry of Health, the National AIDS Control Council (NACC), the National Council for Population and Development (NCPD), and the Kenya Medical Research Institute (KEMRI). Funding for the KDHS was provided by the Government of Kenya with support from the United States Agency for International Development (USAID), the United Nations Population Fund (UNFPA), the United Kingdom Department for International Development (DfID), the World Bank, the Danish International Development Agency (DANIDA), the United Nations Children's Fund (UNICEF), the German Development Bank (KfW), the Clinton Health Access Initiative (CHAI), the World Food Programme (WFP), and the Micronutrient Initiative (MI). ICF International provided technical assistance as well as funding to the project through The DHS Program, a USAID-funded project providing support and technical assistance in the implementation of population and health surveys in countries worldwide.

Republic of Kenya

Additional information about the 2014 KDHS may be obtained from the Kenya National Bureau of Statistics (KNBS), P.O. Box 30266-00100 GPO Nairobi, Kenya; telephone (Nairobi): 3317586/8, 3317612/22, 3317623, 3317651; fax: 3315977; e-mail: directorgeneral@knbs.or.ke, info@knbs.or.ke; website: www.knbs.or.ke.

KNBS KENYA NATIONAL BUREAU OF STATISTICS *Keeping you informed*

Information on The DHS Program may be obtained from ICF International, 530 Gaither Road, Suite 500, Rockville MD, 20850, USA; telephone: 301-407-6500; fax: 301.407.6501; e-mail: info@DHSprogram.com; website: www.DHSprogram.com.



2014 Kenya DHS
Demographic and Health Survey

Taita Taveta


	Taita Taveta	Kenya total
Fertility and Family Planning		
Total fertility rate (number of children per woman)	3.2	3.9
Current use of *any* method of family planning (% of currently married women age 15-49)	68	58
Current use of any modern method of family planning (% of currently married women age 15-49)	61	53
Maternal and Child Health		
Maternity care		
Pregnant women who received antenatal care from a skilled provider[1] (%)	98	96
Pregnant women who received 4+ ANC visits (%)	59	58
Births assisted by a skilled provider[1] (%)	63	62
Births delivered in a health facility (%)	62	61
Child vaccination		
Children 12–23 months fully vaccinated[2] (%)	83	68
Nutrition		
Children under 5 years who are stunted (moderate or severe) (%)	24	26
Children under 5 years who are wasted (moderate or severe) (%)	7	4
Children under 5 years who are underweight (%)	8	11
Children under 5 years who are overweight or obese (%)	4	4
Malaria Prevention and Treatment		
Households with at least one insecticide-treated net[3] (ITN) (%)	80	59
Households with at least one ITN for every 2 persons in the household (universal coverage) (%)	61	34
Children under age 5 who slept under an ITN the night before the survey (%)	82	54
Pregnant women who slept under an ITN the night before the survey (%)	*	51
Pregnant women who received 3+ doses of SP/Fansidar to prevent malaria during pregnancy, at least one dose was received during an ANC visit (IPTp) (%)	32	10
Children under age 5 with fever in the 2 weeks before the survey	30	24
Among children under age 5 with fever, percentage who took any ACT	1	23
HIV/AIDS-related Knowledge		
Women 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	83	77
Men 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	85	85
Women 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	51	53
Men 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	48	46

Note: Figures in parentheses are based on 25-49 unweighted cases; an asterisk indicates that data are based on fewer than 25 unweighted cases and have been suppressed. [1]Skilled provider includes doctor, nurse, or midwife. [2]Fully vaccinated includes BCG, measles, three doses each of DPT and polio vaccine (excluding polio vaccine given at birth) and pneumococcal. [3]An insecticide-treated net (ITN) is (1) a factory-treated net that does not require any further treatment, or (2) a net that has been soaked with insecticide within the past 6 months.

Coast: Key Indicators from the 2014 KDHS


Total Fertility Rate

Average number of live births per woman age 15-49




Use of Modern Methods of Family Planning

Percent of currently married women using a modern method of contraception




Antenatal Care: 4+ visits

Among women with a live birth in the 5 years before the survey, percent with 4+ ANC visits for their last live birth




Delivery in a Health Facility

Percentage of live births delivered in a health facility




Full Vaccination Rate

Percent of children age 12-23 months who have received all basic vaccinations plus the pneumococcal vaccine



ITN Access

Percentage of households with at least one insecticide-treated net (ITN) for every two persons who stayed in the household last night




The 2014 Kenya Demographic and Health Survey (2014 KDHS) was implemented by the Kenya National Bureau of Statistics from May 2014 to October 2014 in partnership with the Ministry of Health, the National AIDS Control Council (NACC), the National Council for Population and Development (NCPD), and the Kenya Medical Research Institute (KEMRI). Funding for the KDHS was provided by the Government of Kenya with support from the United States Agency for International Development (USAID), the United Nations Population Fund (UNFPA), the United Kingdom Department for International Development (DfID), the World Bank, the Danish International Development Agency (DANIDA), the United Nations Children's Fund (UNICEF), the German Development Bank (KfW), the Clinton Health Access Initiative (CHAI), the World Food Programme (WFP), and the Micronutrient Initiative (MI). ICF International provided technical assistance as well as funding to the project through The DHS Program, a USAID-funded project providing support and technical assistance in the implementation of population and health surveys in countries worldwide.

Republic of Kenya

Additional information about the 2014 KDHS may be obtained from the Kenya National Bureau of Statistics (KNBS), P.O. Box 30266-00100 GPO Nairobi, Kenya; telephone (Nairobi): 3317586/8, 3317612/22, 3317623, 3317651; fax: 3315977; e-mail: directorgeneral@knbs.or.ke, info@knbs.or.ke; website: www.knbs.or.ke.

KNBS KENYA NATIONAL BUREAU OF STATISTICS *Keeping you informed*

Information on The DHS Program may be obtained from ICF International, 530 Gaither Road, Suite 500, Rockville MD, 20850, USA; telephone: 301-407-6500; fax: 301.407.6501; e-mail: info@DHSprogram.com; website: www.DHSprogram.com.



2014 Kenya DHS
Demographic and Health Survey

Garissa


	Garissa	Kenya total
Fertility and Family Planning		
Total fertility rate (number of children per woman)	6.1	3.9
Current use of *any* method of family planning (% of currently married women age 15-49)	6	58
Current use of any modern method of family planning (% of currently married women age 15-49)	6	53
Maternal and Child Health		
Maternity care		
Pregnant women who received antenatal care from a skilled provider[1] (%)	87	96
Pregnant women who received 4+ ANC visits (%)	48	58
Births assisted by a skilled provider[1] (%)	40	62
Births delivered in a health facility (%)	37	61
Child vaccination		
Children 12–23 months fully vaccinated[2] (%)	54	68
Nutrition		
Children under 5 years who are stunted (moderate or severe) (%)	16	26
Children under 5 years who are wasted (moderate or severe) (%)	11	4
Children under 5 years who are underweight (%)	13	11
Children under 5 years who are overweight or obese (%)	2	4
Malaria Prevention and Treatment		
Households with at least one insecticide-treated net[3] (ITN) (%)	61	59
Households with at least one ITN for every 2 persons in the household (universal coverage) (%)	34	34
Children under age 5 who slept under an ITN the night before the survey (%)	48	54
Pregnant women who slept under an ITN the night before the survey (%)	52	51
Pregnant women who received 3+ doses of SP/Fansidar to prevent malaria during pregnancy, at least one dose was received during an ANC visit (IPTp) (%)	0.8	10
Children under age 5 with fever in the 2 weeks before the survey	7	24
Among children under age 5 with fever, percentage who took any ACT	(6)	23
HIV/AIDS-related Knowledge		
Women 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	31	77
Men 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	69	85
Women 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	27	53
Men 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	28	46

Note: Figures in parentheses are based on 25-49 unweighted cases; an asterisk indicates that data are based on fewer than 25 unweighted cases and have been suppressed. [1]Skilled provider includes doctor, nurse, or midwife. [2]Fully vaccinated includes BCG, measles, three doses each of DPT and polio vaccine (excluding polio vaccine given at birth) and pneumococcal. [3]An insecticide-treated net (ITN) is (1) a factory-treated net that does not require any further treatment, or (2) a net that has been soaked with insecticide within the past 6 months.

North Eastern: Key Indicators from the 2014 KDHS


Total Fertility Rate

Average number of live births per woman age 15-49




Use of Modern Methods of Family Planning

Percent of currently married women using a modern method of contraception




Antenatal Care: 4+ visits

Among women with a live birth in the 5 years before the survey, percent with 4+ ANC visits for their last live birth




Delivery in a Health Facility

Percentage of live births delivered in a health facility




Full Vaccination Rate

Percent of children age 12-23 months who have received all basic vaccinations plus the pneumococcal vaccine



ITN Access

Percentage of households with at least one insecticide-treated net (ITN) for every two persons who stayed in the household last night




The 2014 Kenya Demographic and Health Survey (2014 KDHS) was implemented by the Kenya National Bureau of Statistics from May 2014 to October 2014 in partnership with the Ministry of Health, the National AIDS Control Council (NACC), the National Council for Population and Development (NCPD), and the Kenya Medical Research Institute (KEMRI). Funding for the KDHS was provided by the Government of Kenya with support from the United States Agency for International Development (USAID), the United Nations Population Fund (UNFPA), the United Kingdom Department for International Development (DfID), the World Bank, the Danish International Development Agency (DANIDA), the United Nations Children's Fund (UNICEF), the German Development Bank (KfW), the Clinton Health Access Initiative (CHAI), the World Food Programme (WFP), and the Micronutrient Initiative (MI). ICF International provided technical assistance as well as funding to the project through The DHS Program, a USAID-funded project providing support and technical assistance in the implementation of population and health surveys in countries worldwide.

Republic of Kenya

Additional information about the 2014 KDHS may be obtained from the Kenya National Bureau of Statistics (KNBS), P.O. Box 30266-00100 GPO Nairobi, Kenya; telephone (Nairobi): 3317586/8, 3317612/22, 3317623, 3317651; fax: 3315977; e-mail: directorgeneral@knbs.or.ke, info@knbs.or.ke; website: www.knbs.or.ke.

Information on The DHS Program may be obtained from ICF International, 530 Gaither Road, Suite 500, Rockville MD, 20850, USA; telephone: 301-407-6500; fax: 301.407.6501; e-mail: info@DHSprogram.com; website: www.DHSprogram.com.



2014 Kenya DHS
Demographic and Health Survey

Wajir


	Wajir	Kenya total
Fertility and Family Planning		
Total fertility rate (number of children per woman)	7.8	3.9
Current use of *any* method of family planning (% of currently married women age 15-49)	2	58
Current use of any modern method of family planning (% of currently married women age 15-49)	2	53
Maternal and Child Health		
Maternity care		
Pregnant women who received antenatal care from a skilled provider[1] (%)	58	96
Pregnant women who received 4+ ANC visits (%)	38	58
Births assisted by a skilled provider[1] (%)	22	62
Births delivered in a health facility (%)	18	61
Child vaccination		
Children 12–23 months fully vaccinated[2] (%)	38	68
Nutrition		
Children under 5 years who are stunted (moderate or severe) (%)	26	26
Children under 5 years who are wasted (moderate or severe) (%)	14	4
Children under 5 years who are underweight (%)	21	11
Children under 5 years who are overweight or obese (%)	0.7	4
Malaria Prevention and Treatment		
Households with at least one insecticide-treated net[3] (ITN) (%)	54	59
Households with at least one ITN for every 2 persons in the household (universal coverage) (%)	18	34
Children under age 5 who slept under an ITN the night before the survey (%)	43	54
Pregnant women who slept under an ITN the night before the survey (%)	49	51
Pregnant women who received 3+ doses of SP/Fansidar to prevent malaria during pregnancy, at least one dose was received during an ANC visit (IPTp) (%)	0.5	10
Children under age 5 with fever in the 2 weeks before the survey	13	24
Among children under age 5 with fever, percentage who took any ACT	0.4	23
HIV/AIDS-related Knowledge		
Women 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	15	77
Men 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	11	85
Women 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	21	53
Men 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	32	46

Note: Figures in parentheses are based on 25-49 unweighted cases; an asterisk indicates that data are based on fewer than 25 unweighted cases and have been suppressed. [1]Skilled provider includes doctor, nurse, or midwife. [2]Fully vaccinated includes BCG, measles, three doses each of DPT and polio vaccine (excluding polio vaccine given at birth) and pneumococcal. [3]An insecticide-treated net (ITN) is (1) a factory-treated net that does not require any further treatment, or (2) a net that has been soaked with insecticide within the past 6 months.

North Eastern: Key Indicators from the 2014 KDHS


Total Fertility Rate

Average number of live births per woman age 15-49




Use of Modern Methods of Family Planning

Percent of currently married women using a modern method of contraception




Antenatal Care: 4+ visits

Among women with a live birth in the 5 years before the survey, percent with 4+ ANC visits for their last live birth




Delivery in a Health Facility

Percentage of live births delivered in a health facility




Full Vaccination Rate

Percent of children age 12-23 months who have received all basic vaccinations plus the pneumococcal vaccine



ITN Access

Percentage of households with at least one insecticide-treated net (ITN) for every two persons who stayed in the household last night




The 2014 Kenya Demographic and Health Survey (2014 KDHS) was implemented by the Kenya National Bureau of Statistics from May 2014 to October 2014 in partnership with the Ministry of Health, the National AIDS Control Council (NACC), the National Council for Population and Development (NCPD), and the Kenya Medical Research Institute (KEMRI). Funding for the KDHS was provided by the Government of Kenya with support from the United States Agency for International Development (USAID), the United Nations Population Fund (UNFPA), the United Kingdom Department for International Development (DfID), the World Bank, the Danish International Development Agency (DANIDA), the United Nations Children's Fund (UNICEF), the German Development Bank (KfW), the Clinton Health Access Initiative (CHAI), the World Food Programme (WFP), and the Micronutrient Initiative (MI). ICF International provided technical assistance as well as funding to the project through The DHS Program, a USAID-funded project providing support and technical assistance in the implementation of population and health surveys in countries worldwide.

Republic of Kenya

Additional information about the 2014 KDHS may be obtained from the Kenya National Bureau of Statistics (KNBS), P.O. Box 30266-00100 GPO Nairobi, Kenya; telephone (Nairobi): 3317586/8, 3317612/22, 3317623, 3317651; fax: 3315977; e-mail: directorgeneral@knbs.or.ke, info@knbs.or.ke; website: www.knbs.or.ke.

Information on The DHS Program may be obtained from ICF International, 530 Gaither Road, Suite 500, Rockville MD, 20850, USA; telephone: 301-407-6500; fax: 301.407.6501; e-mail: info@DHSprogram.com; website: www.DHSprogram.com.



2014 Kenya DHS
Demographic and Health Survey

Mandera


	Mandera	Kenya total
Fertility and Family Planning		
Total fertility rate (number of children per woman)	5.2	3.9
Current use of *any* method of family planning (% of currently married women age 15-49)	2	58
Current use of any modern method of family planning (% of currently married women age 15-49)	2	53
Maternal and Child Health		
Maternity care		
Pregnant women who received antenatal care from a skilled provider[1] (%)	51	96
Pregnant women who received 4+ ANC visits (%)	20	58
Births assisted by a skilled provider[1] (%)	39	62
Births delivered in a health facility (%)	36	61
Child vaccination		
Children 12–23 months fully vaccinated[2] (%)	28	68
Nutrition		
Children under 5 years who are stunted (moderate or severe) (%)	36	26
Children under 5 years who are wasted (moderate or severe) (%)	15	4
Children under 5 years who are underweight (%)	25	11
Children under 5 years who are overweight or obese (%)	6	4
Malaria Prevention and Treatment		
Households with at least one insecticide-treated net[3] (ITN) (%)	29	59
Households with at least one ITN for every 2 persons in the household (universal coverage) (%)	11	34
Children under age 5 who slept under an ITN the night before the survey (%)	26	54
Pregnant women who slept under an ITN the night before the survey (%)	20	51
Pregnant women who received 3+ doses of SP/Fansidar to prevent malaria during pregnancy, at least one dose was received during an ANC visit (IPTp) (%)	3	10
Children under age 5 with fever in the 2 weeks before the survey	5	24
Among children under age 5 with fever, percentage who took any ACT	*	23
HIV/AIDS-related Knowledge		
Women 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	13	77
Men 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	58	85
Women 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	8	53
Men 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	3	46

Note: Figures in parentheses are based on 25-49 unweighted cases; an asterisk indicates that data are based on fewer than 25 unweighted cases and have been suppressed. [1]Skilled provider includes doctor, nurse, or midwife. [2]Fully vaccinated includes BCG, measles, three doses each of DPT and polio vaccine (excluding polio vaccine given at birth) and pneumococcal. [3]An insecticide-treated net (ITN) is (1) a factory-treated net that does not require any further treatment, or (2) a net that has been soaked with insecticide within the past 6 months.

North Eastern: Key Indicators from the 2014 KDHS


Total Fertility Rate

Average number of live births per woman age 15-49




Use of Modern Methods of Family Planning

Percent of currently married women using a modern method of contraception




Antenatal Care: 4+ visits

Among women with a live birth in the 5 years before the survey, percent with 4+ ANC visits for their last live birth




Delivery in a Health Facility

Percentage of live births delivered in a health facility




Full Vaccination Rate

Percent of children age 12-23 months who have received all basic vaccinations plus the pneumococcal vaccine



ITN Access

Percentage of households with at least one insecticide-treated net (ITN) for every two persons who stayed in the household last night




The 2014 Kenya Demographic and Health Survey (2014 KDHS) was implemented by the Kenya National Bureau of Statistics from May 2014 to October 2014 in partnership with the Ministry of Health, the National AIDS Control Council (NACC), the National Council for Population and Development (NCPD), and the Kenya Medical Research Institute (KEMRI). Funding for the KDHS was provided by the Government of Kenya with support from the United States Agency for International Development (USAID), the United Nations Population Fund (UNFPA), the United Kingdom Department for International Development (DfID), the World Bank, the Danish International Development Agency (DANIDA), the United Nations Children's Fund (UNICEF), the German Development Bank (KfW), the Clinton Health Access Initiative (CHAI), the World Food Programme (WFP), and the Micronutrient Initiative (MI). ICF International provided technical assistance as well as funding to the project through The DHS Program, a USAID-funded project providing support and technical assistance in the implementation of population and health surveys in countries worldwide.

Republic of Kenya

Additional information about the 2014 KDHS may be obtained from the Kenya National Bureau of Statistics (KNBS), P.O. Box 30266-00100 GPO Nairobi, Kenya; telephone (Nairobi): 3317586/8, 3317612/22, 3317623, 3317651; fax: 3315977; e-mail: directorgeneral@knbs.or.ke, info@knbs.or.ke; website: www.knbs.or.ke.

Information on The DHS Program may be obtained from ICF International, 530 Gaither Road, Suite 500, Rockville MD, 20850, USA; telephone: 301-407-6500; fax: 301.407.6501; e-mail: info@DHSprogram.com; website: www.DHSprogram.com.



2014 Kenya DHS
Demographic and Health Survey

Marsabit


	Marsabit	Kenya total
Fertility and Family Planning		
Total fertility rate (number of children per woman)	5.0	3.9
Current use of *any* method of family planning (% of currently married women age 15-49)	12	58
Current use of any modern method of family planning (% of currently married women age 15-49)	11	53
Maternal and Child Health		
Maternity care		
Pregnant women who received antenatal care from a skilled provider[1] (%)	76	96
Pregnant women who received 4+ ANC visits (%)	43	58
Births assisted by a skilled provider[1] (%)	26	62
Births delivered in a health facility (%)	26	61
Child vaccination		
Children 12–23 months fully vaccinated[2] (%)	67	68
Nutrition		
Children under 5 years who are stunted (moderate or severe) (%)	27	26
Children under 5 years who are wasted (moderate or severe) (%)	16	4
Children under 5 years who are underweight (%)	30	11
Children under 5 years who are overweight or obese (%)	1	4
Malaria Prevention and Treatment		
Households with at least one insecticide-treated net[3] (ITN) (%)	25	59
Households with at least one ITN for every 2 persons in the household (universal coverage) (%)	7	34
Children under age 5 who slept under an ITN the night before the survey (%)	12	54
Pregnant women who slept under an ITN the night before the survey (%)	8	51
Pregnant women who received 3+ doses of SP/Fansidar to prevent malaria during pregnancy, at least one dose was received during an ANC visit (IPTp) (%)	0.6	10
Children under age 5 with fever in the 2 weeks before the survey	19	24
Among children under age 5 with fever, percentage who took any ACT	10	23
HIV/AIDS-related Knowledge		
Women 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	45	77
Men 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	95	85
Women 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	34	53
Men 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	51	46

Note: Figures in parentheses are based on 25-49 unweighted cases; an asterisk indicates that data are based on fewer than 25 unweighted cases and have been suppressed. [1]Skilled provider includes doctor, nurse, or midwife. [2]Fully vaccinated includes BCG, measles, three doses each of DPT and polio vaccine (excluding polio vaccine given at birth) and pneumococcal. [3]An insecticide-treated net (ITN) is (1) a factory-treated net that does not require any further treatment, or (2) a net that has been soaked with insecticide within the past 6 months.

Eastern: Key Indicators from the 2014 KDHS


Total Fertility Rate

Average number of live births per woman age 15-49




Use of Modern Methods of Family Planning

Percent of currently married women using a modern method of contraception




Antenatal Care: 4+ visits

Among women with a live birth in the 5 years before the survey, percent with 4+ ANC visits for their last live birth




Delivery in a Health Facility

Percentage of live births delivered in a health facility




Full Vaccination Rate

Percent of children age 12-23 months who have received all basic vaccinations plus the pneumococcal vaccine



ITN Access

Percentage of households with at least one insecticide-treated net (ITN) for every two persons who stayed in the household last night




The 2014 Kenya Demographic and Health Survey (2014 KDHS) was implemented by the Kenya National Bureau of Statistics from May 2014 to October 2014 in partnership with the Ministry of Health, the National AIDS Control Council (NACC), the National Council for Population and Development (NCPD), and the Kenya Medical Research Institute (KEMRI). Funding for the KDHS was provided by the Government of Kenya with support from the United States Agency for International Development (USAID), the United Nations Population Fund (UNFPA), the United Kingdom Department for International Development (DfID), the World Bank, the Danish International Development Agency (DANIDA), the United Nations Children's Fund (UNICEF), the German Development Bank (KfW), the Clinton Health Access Initiative (CHAI), the World Food Programme (WFP), and the Micronutrient Initiative (MI). ICF International provided technical assistance as well as funding to the project through The DHS Program, a USAID-funded project providing support and technical assistance in the implementation of population and health surveys in countries worldwide.

Republic of Kenya

Additional information about the 2014 KDHS may be obtained from the Kenya National Bureau of Statistics (KNBS), P.O. Box 30266-00100 GPO Nairobi, Kenya; telephone (Nairobi): 3317586/8, 3317612/22, 3317623, 3317651; fax: 3315977; e-mail: directorgeneral@knbs.or.ke, info@knbs.or.ke; website: www.knbs.or.ke.

KNBS KENYA NATIONAL BUREAU OF STATISTICS *Keeping you informed*

Information on The DHS Program may be obtained from ICF International, 530 Gaither Road, Suite 500, Rockville MD, 20850, USA; telephone: 301-407-6500; fax: 301.407.6501; e-mail: info@DHSprogram.com; website: www.DHSprogram.com.



2014 Kenya DHS
Demographic and Health Survey

Isiolo


	Isiolo	Kenya total
Fertility and Family Planning		
Total fertility rate (number of children per woman)	4.9	3.9
Current use of *any* method of family planning (% of currently married women age 15-49)	27	58
Current use of any modern method of family planning (% of currently married women age 15-49)	26	53
Maternal and Child Health		
Maternity care		
Pregnant women who received antenatal care from a skilled provider[1] (%)	96	96
Pregnant women who received 4+ ANC visits (%)	50	58
Births assisted by a skilled provider[1] (%)	44	62
Births delivered in a health facility (%)	42	61
Child vaccination		
Children 12–23 months fully vaccinated[2] (%)	79	68
Nutrition		
Children under 5 years who are stunted (moderate or severe) (%)	19	26
Children under 5 years who are wasted (moderate or severe) (%)	9	4
Children under 5 years who are underweight (%)	13	11
Children under 5 years who are overweight or obese (%)	2	4
Malaria Prevention and Treatment		
Households with at least one insecticide-treated net[3] (ITN) (%)	63	59
Households with at least one ITN for every 2 persons in the household (universal coverage) (%)	32	34
Children under age 5 who slept under an ITN the night before the survey (%)	58	54
Pregnant women who slept under an ITN the night before the survey (%)	(56)	51
Pregnant women who received 3+ doses of SP/Fansidar to prevent malaria during pregnancy, at least one dose was received during an ANC visit (IPTp) (%)	1	10
Children under age 5 with fever in the 2 weeks before the survey	13	24
Among children under age 5 with fever, percentage who took any ACT	43	23
HIV/AIDS-related Knowledge		
Women 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	74	77
Men 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	71	85
Women 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	46	53
Men 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	41	46

Note: Figures in parentheses are based on 25-49 unweighted cases; an asterisk indicates that data are based on fewer than 25 unweighted cases and have been suppressed. [1]Skilled provider includes doctor, nurse, or midwife. [2]Fully vaccinated includes BCG, measles, three doses each of DPT and polio vaccine (excluding polio vaccine given at birth) and pneumococcal. [3]An insecticide-treated net (ITN) is (1) a factory-treated net that does not require any further treatment, or (2) a net that has been soaked with insecticide within the past 6 months.

Eastern: Key Indicators from the 2014 KDHS


Total Fertility Rate

Average number of live births per woman age 15-49




Use of Modern Methods of Family Planning

Percent of currently married women using a modern method of contraception




Antenatal Care: 4+ visits

Among women with a live birth in the 5 years before the survey, percent with 4+ ANC visits for their last live birth




Delivery in a Health Facility

Percentage of live births delivered in a health facility




Full Vaccination Rate

Percent of children age 12-23 months who have received all basic vaccinations plus the pneumococcal vaccine



ITN Access

Percentage of households with at least one insecticide-treated net (ITN) for every two persons who stayed in the household last night




The 2014 Kenya Demographic and Health Survey (2014 KDHS) was implemented by the Kenya National Bureau of Statistics from May 2014 to October 2014 in partnership with the Ministry of Health, the National AIDS Control Council (NACC), the National Council for Population and Development (NCPD), and the Kenya Medical Research Institute (KEMRI). Funding for the KDHS was provided by the Government of Kenya with support from the United States Agency for International Development (USAID), the United Nations Population Fund (UNFPA), the United Kingdom Department for International Development (DfID), the World Bank, the Danish International Development Agency (DANIDA), the United Nations Children's Fund (UNICEF), the German Development Bank (KfW), the Clinton Health Access Initiative (CHAI), the World Food Programme (WFP), and the Micronutrient Initiative (MI). ICF International provided technical assistance as well as funding to the project through The DHS Program, a USAID-funded project providing support and technical assistance in the implementation of population and health surveys in countries worldwide.

Republic of Kenya

Additional information about the 2014 KDHS may be obtained from the Kenya National Bureau of Statistics (KNBS), P.O. Box 30266-00100 GPO Nairobi, Kenya; telephone (Nairobi): 3317586/8, 3317612/22, 3317623, 3317651; fax: 3315977; e-mail: directorgeneral@knbs.or.ke, info@knbs.or.ke; website: www.knbs.or.ke.

KNBS KENYA NATIONAL BUREAU OF STATISTICS *Keeping you informed*

Information on The DHS Program may be obtained from ICF International, 530 Gaither Road, Suite 500, Rockville MD, 20850, USA; telephone: 301-407-6500; fax: 301.407.6501; e-mail: info@DHSprogram.com; website: www.DHSprogram.com.



2014 Kenya DHS
Demographic and Health Survey

Meru


	Meru	Kenya total
Fertility and Family Planning		
Total fertility rate (number of children per woman)	3.1	3.9
Current use of *any* method of family planning (% of currently married women age 15-49)	78	58
Current use of any modern method of family planning (% of currently married women age 15-49)	73	53
Maternal and Child Health		
Maternity care		
Pregnant women who received antenatal care from a skilled provider[1] (%)	97	96
Pregnant women who received 4+ ANC visits (%)	45	58
Births assisted by a skilled provider[1] (%)	83	62
Births delivered in a health facility (%)	82	61
Child vaccination		
Children 12–23 months fully vaccinated[2] (%)	78	68
Nutrition		
Children under 5 years who are stunted (moderate or severe) (%)	25	26
Children under 5 years who are wasted (moderate or severe) (%)	3	4
Children under 5 years who are underweight (%)	8	11
Children under 5 years who are overweight or obese (%)	5	4
Malaria Prevention and Treatment		
Households with at least one insecticide-treated net[3] (ITN) (%)	54	59
Households with at least one ITN for every 2 persons in the household (universal coverage) (%)	31	34
Children under age 5 who slept under an ITN the night before the survey (%)	59	54
Pregnant women who slept under an ITN the night before the survey (%)	(49)	51
Pregnant women who received 3+ doses of SP/Fansidar to prevent malaria during pregnancy, at least one dose was received during an ANC visit (IPTp) (%)	0	10
Children under age 5 with fever in the 2 weeks before the survey	26	24
Among children under age 5 with fever, percentage who took any ACT	15	23
HIV/AIDS-related Knowledge		
Women 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	79	77
Men 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	86	85
Women 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	45	53
Men 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	38	46

Note: Figures in parentheses are based on 25-49 unweighted cases; an asterisk indicates that data are based on fewer than 25 unweighted cases and have been suppressed. [1]Skilled provider includes doctor, nurse, or midwife. [2]Fully vaccinated includes BCG, measles, three doses each of DPT and polio vaccine (excluding polio vaccine given at birth) and pneumococcal. [3]An insecticide-treated net (ITN) is (1) a factory-treated net that does not require any further treatment, or (2) a net that has been soaked with insecticide within the past 6 months.

Eastern: Key Indicators from the 2014 KDHS


Total Fertility Rate

Average number of live births per woman age 15-49




Use of Modern Methods of Family Planning

Percent of currently married women using a modern method of contraception




Antenatal Care: 4+ visits

Among women with a live birth in the 5 years before the survey, percent with 4+ ANC visits for their last live birth




Delivery in a Health Facility

Percentage of live births delivered in a health facility




Full Vaccination Rate

Percent of children age 12-23 months who have received all basic vaccinations plus the pneumococcal vaccine



ITN Access

Percentage of households with at least one insecticide-treated net (ITN) for every two persons who stayed in the household last night




The 2014 Kenya Demographic and Health Survey (2014 KDHS) was implemented by the Kenya National Bureau of Statistics from May 2014 to October 2014 in partnership with the Ministry of Health, the National AIDS Control Council (NACC), the National Council for Population and Development (NCPD), and the Kenya Medical Research Institute (KEMRI). Funding for the KDHS was provided by the Government of Kenya with support from the United States Agency for International Development (USAID), the United Nations Population Fund (UNFPA), the United Kingdom Department for International Development (DfID), the World Bank, the Danish International Development Agency (DANIDA), the United Nations Children's Fund (UNICEF), the German Development Bank (KfW), the Clinton Health Access Initiative (CHAI), the World Food Programme (WFP), and the Micronutrient Initiative (MI). ICF International provided technical assistance as well as funding to the project through The DHS Program, a USAID-funded project providing support and technical assistance in the implementation of population and health surveys in countries worldwide.

Republic of Kenya

Additional information about the 2014 KDHS may be obtained from the Kenya National Bureau of Statistics (KNBS), P.O. Box 30266-00100 GPO Nairobi, Kenya; telephone (Nairobi): 3317586/8, 3317612/22, 3317623, 3317651; fax: 3315977; e-mail: directorgeneral@knbs.or.ke, info@knbs.or.ke; website: www.knbs.or.ke.

KNBS KENYA NATIONAL BUREAU OF STATISTICS *Keeping you informed*

Information on The DHS Program may be obtained from ICF International, 530 Gaither Road, Suite 500, Rockville MD, 20850, USA; telephone: 301-407-6500; fax: 301.407.6501; e-mail: info@DHSprogram.com; website: www.DHSprogram.com.



2014 Kenya DHS
Demographic and Health Survey

Tharaka-Nithi


	Tharaka-Nithi	Kenya total
Fertility and Family Planning		
Total fertility rate (number of children per woman)	3.4	3.9
Current use of *any* method of family planning (% of currently married women age 15-49)	74	58
Current use of any modern method of family planning (% of currently married women age 15-49)	67	53
Maternal and Child Health		
Maternity care		
Pregnant women who received antenatal care from a skilled provider[1] (%)	98	96
Pregnant women who received 4+ ANC visits (%)	56	58
Births assisted by a skilled provider[1] (%)	77	62
Births delivered in a health facility (%)	78	61
Child vaccination		
Children 12–23 months fully vaccinated[2] (%)	91	68
Nutrition		
Children under 5 years who are stunted (moderate or severe) (%)	33	26
Children under 5 years who are wasted (moderate or severe) (%)	3	4
Children under 5 years who are underweight (%)	11	11
Children under 5 years who are overweight or obese (%)	4	4
Malaria Prevention and Treatment		
Households with at least one insecticide-treated net[3] (ITN) (%)	67	59
Households with at least one ITN for every 2 persons in the household (universal coverage) (%)	46	34
Children under age 5 who slept under an ITN the night before the survey (%)	62	54
Pregnant women who slept under an ITN the night before the survey (%)	*	51
Pregnant women who received 3+ doses of SP/Fansidar to prevent malaria during pregnancy, at least one dose was received during an ANC visit (IPTp) (%)	7	10
Children under age 5 with fever in the 2 weeks before the survey	28	24
Among children under age 5 with fever, percentage who took any ACT	10	23
HIV/AIDS-related Knowledge		
Women 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	74	77
Men 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	81	85
Women 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	52	53
Men 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	43	46

Note: Figures in parentheses are based on 25-49 unweighted cases; an asterisk indicates that data are based on fewer than 25 unweighted cases and have been suppressed. [1]Skilled provider includes doctor, nurse, or midwife. [2]Fully vaccinated includes BCG, measles, three doses each of DPT and polio vaccine (excluding polio vaccine given at birth) and pneumococcal. [3]An insecticide-treated net (ITN) is (1) a factory-treated net that does not require any further treatment, or (2) a net that has been soaked with insecticide within the past 6 months.

Eastern: Key Indicators from the 2014 KDHS


Total Fertility Rate

Average number of live births per woman age 15-49




Use of Modern Methods of Family Planning

Percent of currently married women using a modern method of contraception




Antenatal Care: 4+ visits

Among women with a live birth in the 5 years before the survey, percent with 4+ ANC visits for their last live birth




Delivery in a Health Facility

Percentage of live births delivered in a health facility




Full Vaccination Rate

Percent of children age 12-23 months who have received all basic vaccinations plus the pneumococcal vaccine



ITN Access

Percentage of households with at least one insecticide-treated net (ITN) for every two persons who stayed in the household last night




The 2014 Kenya Demographic and Health Survey (2014 KDHS) was implemented by the Kenya National Bureau of Statistics from May 2014 to October 2014 in partnership with the Ministry of Health, the National AIDS Control Council (NACC), the National Council for Population and Development (NCPD), and the Kenya Medical Research Institute (KEMRI). Funding for the KDHS was provided by the Government of Kenya with support from the United States Agency for International Development (USAID), the United Nations Population Fund (UNFPA), the United Kingdom Department for International Development (DfID), the World Bank, the Danish International Development Agency (DANIDA), the United Nations Children's Fund (UNICEF), the German Development Bank (KfW), the Clinton Health Access Initiative (CHAI), the World Food Programme (WFP), and the Micronutrient Initiative (MI). ICF International provided technical assistance as well as funding to the project through The DHS Program, a USAID-funded project providing support and technical assistance in the implementation of population and health surveys in countries worldwide.

Republic of Kenya

Additional information about the 2014 KDHS may be obtained from the Kenya National Bureau of Statistics (KNBS), P.O. Box 30266-00100 GPO Nairobi, Kenya; telephone (Nairobi): 3317586/8, 3317612/22, 3317623, 3317651; fax: 3315977; e-mail: directorgeneral@knbs.or.ke, info@knbs.or.ke; website: www.knbs.or.ke.

KNBS KENYA NATIONAL BUREAU OF STATISTICS *Keeping you informed*

Information on The DHS Program may be obtained from ICF International, 530 Gaither Road, Suite 500, Rockville MD, 20850, USA; telephone: 301-407-6500; fax: 301.407.6501; e-mail: info@DHSprogram.com; website: www.DHSprogram.com.



2014 Kenya DHS
Demographic and Health Survey

Embu


	Embu	Kenya total
Fertility and Family Planning		
Total fertility rate (number of children per woman)	3.1	3.9
Current use of *any* method of family planning (% of currently married women age 15-49)	71	58
Current use of any modern method of family planning (% of currently married women age 15-49)	67	53
Maternal and Child Health		
Maternity care		
Pregnant women who received antenatal care from a skilled provider[1] (%)	99	96
Pregnant women who received 4+ ANC visits (%)	56	58
Births assisted by a skilled provider[1] (%)	82	62
Births delivered in a health facility (%)	82	61
Child vaccination		
Children 12–23 months fully vaccinated[2] (%)	83	68
Nutrition		
Children under 5 years who are stunted (moderate or severe) (%)	27	26
Children under 5 years who are wasted (moderate or severe) (%)	3	4
Children under 5 years who are underweight (%)	11	11
Children under 5 years who are overweight or obese (%)	4	4
Malaria Prevention and Treatment		
Households with at least one insecticide-treated net[3] (ITN) (%)	56	59
Households with at least one ITN for every 2 persons in the household (universal coverage) (%)	35	34
Children under age 5 who slept under an ITN the night before the survey (%)	59	54
Pregnant women who slept under an ITN the night before the survey (%)	*	51
Pregnant women who received 3+ doses of SP/Fansidar to prevent malaria during pregnancy, at least one dose was received during an ANC visit (IPTp) (%)	0	10
Children under age 5 with fever in the 2 weeks before the survey	14	24
Among children under age 5 with fever, percentage who took any ACT	(22)	23
HIV/AIDS-related Knowledge		
Women 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	78	77
Men 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	63	85
Women 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	51	53
Men 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	36	46

Note: Figures in parentheses are based on 25-49 unweighted cases; an asterisk indicates that data are based on fewer than 25 unweighted cases and have been suppressed. [1]Skilled provider includes doctor, nurse, or midwife. [2]Fully vaccinated includes BCG, measles, three doses each of DPT and polio vaccine (excluding polio vaccine given at birth) and pneumococcal. [3]An insecticide-treated net (ITN) is (1) a factory-treated net that does not require any further treatment, or (2) a net that has been soaked with insecticide within the past 6 months.

Eastern: Key Indicators from the 2014 KDHS


Total Fertility Rate

Average number of live births per woman age 15-49




Use of Modern Methods of Family Planning

Percent of currently married women using a modern method of contraception




Antenatal Care: 4+ visits

Among women with a live birth in the 5 years before the survey, percent with 4+ ANC visits for their last live birth




Delivery in a Health Facility

Percentage of live births delivered in a health facility




Full Vaccination Rate

Percent of children age 12-23 months who have received all basic vaccinations plus the pneumococcal vaccine



ITN Access

Percentage of households with at least one insecticide-treated net (ITN) for every two persons who stayed in the household last night




The 2014 Kenya Demographic and Health Survey (2014 KDHS) was implemented by the Kenya National Bureau of Statistics from May 2014 to October 2014 in partnership with the Ministry of Health, the National AIDS Control Council (NACC), the National Council for Population and Development (NCPD), and the Kenya Medical Research Institute (KEMRI). Funding for the KDHS was provided by the Government of Kenya with support from the United States Agency for International Development (USAID), the United Nations Population Fund (UNFPA), the United Kingdom Department for International Development (DfID), the World Bank, the Danish International Development Agency (DANIDA), the United Nations Children's Fund (UNICEF), the German Development Bank (KfW), the Clinton Health Access Initiative (CHAI), the World Food Programme (WFP), and the Micronutrient Initiative (MI). ICF International provided technical assistance as well as funding to the project through The DHS Program, a USAID-funded project providing support and technical assistance in the implementation of population and health surveys in countries worldwide.

Republic of Kenya

Additional information about the 2014 KDHS may be obtained from the Kenya National Bureau of Statistics (KNBS), P.O. Box 30266-00100 GPO Nairobi, Kenya; telephone (Nairobi): 3317586/8, 3317612/22, 3317623, 3317651; fax: 3315977; e-mail: directorgeneral@knbs.or.ke, info@knbs.or.ke; website: www.knbs.or.ke.

KNBS KENYA NATIONAL BUREAU OF STATISTICS *Keeping you informed*

Information on The DHS Program may be obtained from ICF International, 530 Gaither Road, Suite 500, Rockville MD, 20850, USA; telephone: 301-407-6500; fax: 301.407.6501; e-mail: info@DHSprogram.com; website: www.DHSprogram.com.



KITUI

2014 KENYA DHS
Demographic and Health Survey


	Kitui	Kenya total
Fertility and Family Planning		
Total fertility rate (number of children per woman)	3.9	3.9
Current use of *any* method of family planning (% of currently married women age 15-49)	57	58
Current use of any modern method of family planning (% of currently married women age 15-49)	55	53
Maternal and Child Health		
Maternity care		
Pregnant women who received antenatal care from a skilled provider[1] (%)	98	96
Pregnant women who received 4+ ANC visits (%)	62	58
Births assisted by a skilled provider[1] (%)	46	62
Births delivered in a health facility (%)	46	61
Child vaccination		
Children 12–23 months fully vaccinated[2] (%)	53	68
Nutrition		
Children under 5 years who are stunted (moderate or severe) (%)	46	26
Children under 5 years who are wasted (moderate or severe) (%)	3	4
Children under 5 years who are underweight (%)	20	11
Children under 5 years who are overweight or obese (%)	3	4
Malaria Prevention and Treatment		
Households with at least one insecticide-treated net[3] (ITN) (%)	61	59
Households with at least one ITN for every 2 persons in the household (universal coverage) (%)	29	34
Children under age 5 who slept under an ITN the night before the survey (%)	40	54
Pregnant women who slept under an ITN the night before the survey (%)	(47)	51
Pregnant women who received 3+ doses of SP/Fansidar to prevent malaria during pregnancy, at least one dose was received during an ANC visit (IPTp) (%)	16	10
Children under age 5 with fever in the 2 weeks before the survey	17	24
Among children under age 5 with fever, percentage who took any ACT	4	23
HIV/AIDS-related Knowledge		
Women 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	50	77
Men 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	82	85
Women 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	56	53
Men 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	33	46

Note: Figures in parentheses are based on 25-49 unweighted cases; an asterisk indicates that data are based on fewer than 25 unweighted cases and have been suppressed. [1]Skilled provider includes doctor, nurse, or midwife. [2]Fully vaccinated includes BCG, measles, three doses each of DPT and polio vaccine (excluding polio vaccine given at birth) and pneumococcal. [3]An insecticide-treated net (ITN) is (1) a factory-treated net that does not require any further treatment, or (2) a net that has been soaked with insecticide within the past 6 months.

Eastern: Key Indicators from the 2014 KDHS


Total Fertility Rate

Average number of live births per woman age 15-49




Use of Modern Methods of Family Planning

Percent of currently married women using a modern method of contraception




Antenatal Care: 4+ visits

Among women with a live birth in the 5 years before the survey, percent with 4+ ANC visits for their last live birth




Delivery in a Health Facility

Percentage of live births delivered in a health facility




Full Vaccination Rate

Percent of children age 12-23 months who have received all basic vaccinations plus the pneumococcal vaccine



ITN Access

Percentage of households with at least one insecticide-treated net (ITN) for every two persons who stayed in the household last night




The 2014 Kenya Demographic and Health Survey (2014 KDHS) was implemented by the Kenya National Bureau of Statistics from May 2014 to October 2014 in partnership with the Ministry of Health, the National AIDS Control Council (NACC), the National Council for Population and Development (NCPD), and the Kenya Medical Research Institute (KEMRI). Funding for the KDHS was provided by the Government of Kenya with support from the United States Agency for International Development (USAID), the United Nations Population Fund (UNFPA), the United Kingdom Department for International Development (DfID), the World Bank, the Danish International Development Agency (DANIDA), the United Nations Children's Fund (UNICEF), the German Development Bank (KfW), the Clinton Health Access Initiative (CHAI), the World Food Programme (WFP), and the Micronutrient Initiative (MI). ICF International provided technical assistance as well as funding to the project through The DHS Program, a USAID-funded project providing support and technical assistance in the implementation of population and health surveys in countries worldwide.

Republic of Kenya

Additional information about the 2014 KDHS may be obtained from the Kenya National Bureau of Statistics (KNBS), P.O. Box 30266-00100 GPO Nairobi, Kenya; telephone (Nairobi): 3317586/8, 3317612/22, 3317623, 3317651; fax: 3315977; e-mail: directorgeneral@knbs.or.ke, info@knbs.or.ke; website: www.knbs.or.ke.

KNBS KENYA NATIONAL BUREAU OF STATISTICS *Keeping you informed*

Information on The DHS Program may be obtained from ICF International, 530 Gaither Road, Suite 500, Rockville MD, 20850, USA; telephone: 301-407-6500; fax: 301.407.6501; e-mail: info@DHSprogram.com; website: www.DHSprogram.com.



2014 Kenya DHS
Demographic and Health Survey

Machakos


	Machakos	Kenya total
Fertility and Family Planning		
Total fertility rate (number of children per woman)	3.4	3.9
Current use of *any* method of family planning (% of currently married women age 15-49)	76	58
Current use of any modern method of family planning (% of currently married women age 15-49)	68	53
Maternal and Child Health		
Maternity care		
Pregnant women who received antenatal care from a skilled provider[1] (%)	99	96
Pregnant women who received 4+ ANC visits (%)	61	58
Births assisted by a skilled provider[1] (%)	63	62
Births delivered in a health facility (%)	63	61
Child vaccination		
Children 12–23 months fully vaccinated[2] (%)	87	68
Nutrition		
Children under 5 years who are stunted (moderate or severe) (%)	27	26
Children under 5 years who are wasted (moderate or severe) (%)	7	4
Children under 5 years who are underweight (%)	8	11
Children under 5 years who are overweight or obese (%)	6	4
Malaria Prevention and Treatment		
Households with at least one insecticide-treated net[3] (ITN) (%)	56	59
Households with at least one ITN for every 2 persons in the household (universal coverage) (%)	32	34
Children under age 5 who slept under an ITN the night before the survey (%)	63	54
Pregnant women who slept under an ITN the night before the survey (%)	*	51
Pregnant women who received 3+ doses of SP/Fansidar to prevent malaria during pregnancy, at least one dose was received during an ANC visit (IPTp) (%)	3	10
Children under age 5 with fever in the 2 weeks before the survey	14	24
Among children under age 5 with fever, percentage who took any ACT	(11)	23
HIV/AIDS-related Knowledge		
Women 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	75	77
Men 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	83	85
Women 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	55	53
Men 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	46	46

Note: Figures in parentheses are based on 25-49 unweighted cases; an asterisk indicates that data are based on fewer than 25 unweighted cases and have been suppressed. [1]Skilled provider includes doctor, nurse, or midwife. [2]Fully vaccinated includes BCG, measles, three doses each of DPT and polio vaccine (excluding polio vaccine given at birth) and pneumococcal. [3]An insecticide-treated net (ITN) is (1) a factory-treated net that does not require any further treatment, or (2) a net that has been soaked with insecticide within the past 6 months.

Eastern: Key Indicators from the 2014 KDHS


Total Fertility Rate

Average number of live births per woman age 15-49




Use of Modern Methods of Family Planning

Percent of currently married women using a modern method of contraception




Antenatal Care: 4+ visits

Among women with a live birth in the 5 years before the survey, percent with 4+ ANC visits for their last live birth




Delivery in a Health Facility

Percentage of live births delivered in a health facility




Full Vaccination Rate

Percent of children age 12-23 months who have received all basic vaccinations plus the pneumococcal vaccine



ITN Access

Percentage of households with at least one insecticide-treated net (ITN) for every two persons who stayed in the household last night




The 2014 Kenya Demographic and Health Survey (2014 KDHS) was implemented by the Kenya National Bureau of Statistics from May 2014 to October 2014 in partnership with the Ministry of Health, the National AIDS Control Council (NACC), the National Council for Population and Development (NCPD), and the Kenya Medical Research Institute (KEMRI). Funding for the KDHS was provided by the Government of Kenya with support from the United States Agency for International Development (USAID), the United Nations Population Fund (UNFPA), the United Kingdom Department for International Development (DfID), the World Bank, the Danish International Development Agency (DANIDA), the United Nations Children's Fund (UNICEF), the German Development Bank (KfW), the Clinton Health Access Initiative (CHAI), the World Food Programme (WFP), and the Micronutrient Initiative (MI). ICF International provided technical assistance as well as funding to the project through The DHS Program, a USAID-funded project providing support and technical assistance in the implementation of population and health surveys in countries worldwide.

Republic of Kenya

Additional information about the 2014 KDHS may be obtained from the Kenya National Bureau of Statistics (KNBS), P.O. Box 30266-00100 GPO Nairobi, Kenya; telephone (Nairobi): 3317586/8, 3317612/22, 3317623, 3317651; fax: 3315977; e-mail: directorgeneral@knbs.or.ke, info@knbs.or.ke; website: www.knbs.or.ke.

KNBS KENYA NATIONAL BUREAU OF STATISTICS *Keeping you informed*

Information on The DHS Program may be obtained from ICF International, 530 Gaither Road, Suite 500, Rockville MD, 20850, USA; telephone: 301-407-6500; fax: 301.407.6501; e-mail: info@DHSprogram.com; website: www.DHSprogram.com.



2014 Kenya DHS
Demographic and Health Survey

Makueni


	Makueni	Kenya total
Fertility and Family Planning		
Total fertility rate (number of children per woman)	3.3	3.9
Current use of *any* method of family planning (% of currently married women age 15-49)	80	58
Current use of any modern method of family planning (% of currently married women age 15-49)	65	53
Maternal and Child Health		
Maternity care		
Pregnant women who received antenatal care from a skilled provider[1] (%)	98	96
Pregnant women who received 4+ ANC visits (%)	66	58
Births assisted by a skilled provider[1] (%)	55	62
Births delivered in a health facility (%)	53	61
Child vaccination		
Children 12–23 months fully vaccinated[2] (%)	90	68
Nutrition		
Children under 5 years who are stunted (moderate or severe) (%)	25	26
Children under 5 years who are wasted (moderate or severe) (%)	2	4
Children under 5 years who are underweight (%)	10	11
Children under 5 years who are overweight or obese (%)	5	4
Malaria Prevention and Treatment		
Households with at least one insecticide-treated net[3] (ITN) (%)	55	59
Households with at least one ITN for every 2 persons in the household (universal coverage) (%)	28	34
Children under age 5 who slept under an ITN the night before the survey (%)	51	54
Pregnant women who slept under an ITN the night before the survey (%)	(48)	51
Pregnant women who received 3+ doses of SP/Fansidar to prevent malaria during pregnancy, at least one dose was received during an ANC visit (IPTp) (%)	7	10
Children under age 5 with fever in the 2 weeks before the survey	14	24
Among children under age 5 with fever, percentage who took any ACT	8	23
HIV/AIDS-related Knowledge		
Women 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	77	77
Men 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	92	85
Women 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	52	53
Men 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	40	46

Note: Figures in parentheses are based on 25-49 unweighted cases; an asterisk indicates that data are based on fewer than 25 unweighted cases and have been suppressed. [1]Skilled provider includes doctor, nurse, or midwife. [2]Fully vaccinated includes BCG, measles, three doses each of DPT and polio vaccine (excluding polio vaccine given at birth) and pneumococcal. [3]An insecticide-treated net (ITN) is (1) a factory-treated net that does not require any further treatment, or (2) a net that has been soaked with insecticide within the past 6 months.

Eastern: Key Indicators from the 2014 KDHS


Total Fertility Rate

Average number of live births per woman age 15-49




Use of Modern Methods of Family Planning

Percent of currently married women using a modern method of contraception




Antenatal Care: 4+ visits

Among women with a live birth in the 5 years before the survey, percent with 4+ ANC visits for their last live birth




Delivery in a Health Facility

Percentage of live births delivered in a health facility




Full Vaccination Rate

Percent of children age 12-23 months who have received all basic vaccinations plus the pneumococcal vaccine



ITN Access

Percentage of households with at least one insecticide-treated net (ITN) for every two persons who stayed in the household last night



The 2014 Kenya Demographic and Health Survey (2014 KDHS) was implemented by the Kenya National Bureau of Statistics from May 2014 to October 2014 in partnership with the Ministry of Health, the National AIDS Control Council (NACC), the National Council for Population and Development (NCPD), and the Kenya Medical Research Institute (KEMRI). Funding for the KDHS was provided by the Government of Kenya with support from the United States Agency for International Development (USAID), the United Nations Population Fund (UNFPA), the United Kingdom Department for International Development (DfID), the World Bank, the Danish International Development Agency (DANIDA), the United Nations Children's Fund (UNICEF), the German Development Bank (KfW), the Clinton Health Access Initiative (CHAI), the World Food Programme (WFP), and the Micronutrient Initiative (MI). ICF International provided technical assistance as well as funding to the project through The DHS Program, a USAID-funded project providing support and technical assistance in the implementation of population and health surveys in countries worldwide.

Republic of Kenya

Additional information about the 2014 KDHS may be obtained from the Kenya National Bureau of Statistics (KNBS), P.O. Box 30266-00100 GPO Nairobi, Kenya; telephone (Nairobi): 3317586/8, 3317612/22, 3317623, 3317651; fax: 3315977; e-mail: directorgeneral@knbs.or.ke, info@knbs.or.ke; website: www.knbs.or.ke.

KNBS KENYA NATIONAL BUREAU OF STATISTICS *Keeping you informed*

Information on The DHS Program may be obtained from ICF International, 530 Gaither Road, Suite 500, Rockville MD, 20850, USA; telephone: 301-407-6500; fax: 301.407.6501; e-mail: info@DHSprogram.com; website: www.DHSprogram.com.



2014 KENYA DHS
Demographic and Health Survey

NYANDARUA


	Nyandarua	Kenya total
Fertility and Family Planning		
Total fertility rate (number of children per woman)	3.5	3.9
Current use of *any* method of family planning (% of currently married women age 15-49)	66	58
Current use of any modern method of family planning (% of currently married women age 15-49)	60	53
Maternal and Child Health		
Maternity care		
Pregnant women who received antenatal care from a skilled provider[1] (%)	97	96
Pregnant women who received 4+ ANC visits (%)	58	58
Births assisted by a skilled provider[1] (%)	85	62
Births delivered in a health facility (%)	86	61
Child vaccination		
Children 12–23 months fully vaccinated[2] (%)	78	68
Nutrition		
Children under 5 years who are stunted (moderate or severe) (%)	29	26
Children under 5 years who are wasted (moderate or severe) (%)	2	4
Children under 5 years who are underweight (%)	7	11
Children under 5 years who are overweight or obese (%)	7	4
Malaria Prevention and Treatment		
Households with at least one insecticide-treated net[3] (ITN) (%)	13	59
Households with at least one ITN for every 2 persons in the household (universal coverage) (%)	7	34
Children under age 5 who slept under an ITN the night before the survey (%)	12	54
Pregnant women who slept under an ITN the night before the survey (%)	(7)	51
Pregnant women who received 3+ doses of SP/Fansidar to prevent malaria during pregnancy, at least one dose was received during an ANC visit (IPTp) (%)	2	10
Children under age 5 with fever in the 2 weeks before the survey	17	24
Among children under age 5 with fever, percentage who took any ACT	5	23
HIV/AIDS-related Knowledge		
Women 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	80	77
Men 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	85	85
Women 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	55	53
Men 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	40	46

Note: Figures in parentheses are based on 25-49 unweighted cases; an asterisk indicates that data are based on fewer than 25 unweighted cases and have been suppressed. [1]Skilled provider includes doctor, nurse, or midwife. [2]Fully vaccinated includes BCG, measles, three doses each of DPT and polio vaccine (excluding polio vaccine given at birth) and pneumococcal. [3]An insecticide-treated net (ITN) is (1) a factory-treated net that does not require any further treatment, or (2) a net that has been soaked with insecticide within the past 6 months.

Central and Nairobi: Key Indicators from the 2014 KDHS


Total Fertility Rate

Average number of live births per woman age 15-49




Use of Modern Methods of Family Planning

Percent of currently married women using a modern method of contraception




Antenatal Care: 4+ visits

Among women with a live birth in the 5 years before the survey, percent with 4+ ANC visits for their last live birth




Delivery in a Health Facility

Percentage of live births delivered in a health facility




Full Vaccination Rate

Percent of children age 12-23 months who have received all basic vaccinations plus the pneumococcal vaccine



ITN Access

Percentage of households with at least one insecticide-treated net (ITN) for every two persons who stayed in the household last night



The 2014 Kenya Demographic and Health Survey (2014 KDHS) was implemented by the Kenya National Bureau of Statistics from May 2014 to October 2014 in partnership with the Ministry of Health, the National AIDS Control Council (NACC), the National Council for Population and Development (NCPD), and the Kenya Medical Research Institute (KEMRI). Funding for the KDHS was provided by the Government of Kenya with support from the United States Agency for International Development (USAID), the United Nations Population Fund (UNFPA), the United Kingdom Department for International Development (DfID), the World Bank, the Danish International Development Agency (DANIDA), the United Nations Children's Fund (UNICEF), the German Development Bank (KfW), the Clinton Health Access Initiative (CHAI), the World Food Programme (WFP), and the Micronutrient Initiative (MI). ICF International provided technical assistance as well as funding to the project through The DHS Program, a USAID-funded project providing support and technical assistance in the implementation of population and health surveys in countries worldwide.

Republic of Kenya

Additional information about the 2014 KDHS may be obtained from the Kenya National Bureau of Statistics (KNBS), P.O. Box 30266-00100 GPO Nairobi, Kenya; telephone (Nairobi): 3317586/8, 3317612/22, 3317623, 3317651; fax: 3315977; e-mail: directorgeneral@knbs.or.ke, info@knbs.or.ke; website: www.knbs.or.ke.

KNBS KENYA NATIONAL BUREAU OF STATISTICS *Keeping you informed*

Information on The DHS Program may be obtained from ICF International, 530 Gaither Road, Suite 500, Rockville MD, 20850, USA; telephone: 301-407-6500; fax: 301.407.6501; e-mail: info@DHSprogram.com; website: www.DHSprogram.com.



2014 Kenya DHS
Demographic and Health Survey

Nyeri


	Nyeri	Kenya total
Fertility and Family Planning		
Total fertility rate (number of children per woman)	2.7	3.9
Current use of *any* method of family planning (% of currently married women age 15-49)	73	58
Current use of any modern method of family planning (% of currently married women age 15-49)	67	53
Maternal and Child Health		
Maternity care		
Pregnant women who received antenatal care from a skilled provider[1] (%)	97	96
Pregnant women who received 4+ ANC visits (%)	61	58
Births assisted by a skilled provider[1] (%)	88	62
Births delivered in a health facility (%)	89	61
Child vaccination		
Children 12–23 months fully vaccinated[2] (%)	78	68
Nutrition		
Children under 5 years who are stunted (moderate or severe) (%)	15	26
Children under 5 years who are wasted (moderate or severe) (%)	2	4
Children under 5 years who are underweight (%)	3	11
Children under 5 years who are overweight or obese (%)	6	4
Malaria Prevention and Treatment		
Households with at least one insecticide-treated net[3] (ITN) (%)	20	59
Households with at least one ITN for every 2 persons in the household (universal coverage) (%)	12	34
Children under age 5 who slept under an ITN the night before the survey (%)	17	54
Pregnant women who slept under an ITN the night before the survey (%)	(14)	51
Pregnant women who received 3+ doses of SP/Fansidar to prevent malaria during pregnancy, at least one dose was received during an ANC visit (IPTp) (%)	2	10
Children under age 5 with fever in the 2 weeks before the survey	14	24
Among children under age 5 with fever, percentage who took any ACT	(0)	23
HIV/AIDS-related Knowledge		
Women 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	77	77
Men 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	85	85
Women 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	55	53
Men 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	47	46

Note: Figures in parentheses are based on 25-49 unweighted cases; an asterisk indicates that data are based on fewer than 25 unweighted cases and have been suppressed. [1]Skilled provider includes doctor, nurse, or midwife. [2]Fully vaccinated includes BCG, measles, three doses each of DPT and polio vaccine (excluding polio vaccine given at birth) and pneumococcal. [3]An insecticide-treated net (ITN) is (1) a factory-treated net that does not require any further treatment, or (2) a net that has been soaked with insecticide within the past 6 months.

Central and Nairobi: Key Indicators from the 2014 KDHS


Total Fertility Rate

Average number of live births per woman age 15-49




Use of Modern Methods of Family Planning

Percent of currently married women using a modern method of contraception




Antenatal Care: 4+ visits

Among women with a live birth in the 5 years before the survey, percent with 4+ ANC visits for their last live birth




Delivery in a Health Facility

Percentage of live births delivered in a health facility




Full Vaccination Rate

Percent of children age 12-23 months who have received all basic vaccinations plus the pneumococcal vaccine



ITN Access

Percentage of households with at least one insecticide-treated net (ITN) for every two persons who stayed in the household last night



The 2014 Kenya Demographic and Health Survey (2014 KDHS) was implemented by the Kenya National Bureau of Statistics from May 2014 to October 2014 in partnership with the Ministry of Health, the National AIDS Control Council (NACC), the National Council for Population and Development (NCPD), and the Kenya Medical Research Institute (KEMRI). Funding for the KDHS was provided by the Government of Kenya with support from the United States Agency for International Development (USAID), the United Nations Population Fund (UNFPA), the United Kingdom Department for International Development (DfID), the World Bank, the Danish International Development Agency (DANIDA), the United Nations Children's Fund (UNICEF), the German Development Bank (KfW), the Clinton Health Access Initiative (CHAI), the World Food Programme (WFP), and the Micronutrient Initiative (MI). ICF International provided technical assistance as well as funding to the project through The DHS Program, a USAID-funded project providing support and technical assistance in the implementation of population and health surveys in countries worldwide.

Republic of Kenya

Additional information about the 2014 KDHS may be obtained from the Kenya National Bureau of Statistics (KNBS), P.O. Box 30266-00100 GPO Nairobi, Kenya; telephone (Nairobi): 3317586/8, 3317612/22, 3317623, 3317651; fax: 3315977; e-mail: directorgeneral@knbs.or.ke, info@knbs.or.ke; website: www.knbs.or.ke.

KNBS KENYA NATIONAL BUREAU OF STATISTICS *Keeping you informed*

Information on The DHS Program may be obtained from ICF International, 530 Gaither Road, Suite 500, Rockville MD, 20850, USA; telephone: 301-407-6500; fax: 301.407.6501; e-mail: info@DHSprogram.com; website: www.DHSprogram.com.



2014 KENYA DHS
Demographic and Health Survey

KIRINYAGA


	Kirinyaga	Kenya total
Fertility and Family Planning		
Total fertility rate (number of children per woman)	2.3	3.9
Current use of *any* method of family planning (% of currently married women age 15-49)	81	58
Current use of any modern method of family planning (% of currently married women age 15-49)	76	53
Maternal and Child Health		
Maternity care		
Pregnant women who received antenatal care from a skilled provider[1] (%)	96	96
Pregnant women who received 4+ ANC visits (%)	60	58
Births assisted by a skilled provider[1] (%)	92	62
Births delivered in a health facility (%)	93	61
Child vaccination		
Children 12–23 months fully vaccinated[2] (%)	(55)	68
Nutrition		
Children under 5 years who are stunted (moderate or severe) (%)	17	26
Children under 5 years who are wasted (moderate or severe) (%)	4	4
Children under 5 years who are underweight (%)	8	11
Children under 5 years who are overweight or obese (%)	5	4
Malaria Prevention and Treatment		
Households with at least one insecticide-treated net[3] (ITN) (%)	69	59
Households with at least one ITN for every 2 persons in the household (universal coverage) (%)	54	34
Children under age 5 who slept under an ITN the night before the survey (%)	79	54
Pregnant women who slept under an ITN the night before the survey (%)	(63)	51
Pregnant women who received 3+ doses of SP/Fansidar to prevent malaria during pregnancy, at least one dose was received during an ANC visit (IPTp) (%)	4	10
Children under age 5 with fever in the 2 weeks before the survey	21	24
Among children under age 5 with fever, percentage who took any ACT	(19)	23
HIV/AIDS-related Knowledge		
Women 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	91	77
Men 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	86	85
Women 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	57	53
Men 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	36	46

Note: Figures in parentheses are based on 25-49 unweighted cases; an asterisk indicates that data are based on fewer than 25 unweighted cases and have been suppressed. [1]Skilled provider includes doctor, nurse, or midwife. [2]Fully vaccinated includes BCG, measles, three doses each of DPT and polio vaccine (excluding polio vaccine given at birth) and pneumococcal. [3]An insecticide-treated net (ITN) is (1) a factory-treated net that does not require any further treatment, or (2) a net that has been soaked with insecticide within the past 6 months.

Central and Nairobi: Key Indicators from the 2014 KDHS


Total Fertility Rate

Average number of live births per woman age 15-49




Use of Modern Methods of Family Planning

Percent of currently married women using a modern method of contraception




Antenatal Care: 4+ visits

Among women with a live birth in the 5 years before the survey, percent with 4+ ANC visits for their last live birth




Delivery in a Health Facility

Percentage of live births delivered in a health facility




Full Vaccination Rate

Percent of children age 12-23 months who have received all basic vaccinations plus the pneumococcal vaccine



ITN Access

Percentage of households with at least one insecticide-treated net (ITN) for every two persons who stayed in the household last night



The 2014 Kenya Demographic and Health Survey (2014 KDHS) was implemented by the Kenya National Bureau of Statistics from May 2014 to October 2014 in partnership with the Ministry of Health, the National AIDS Control Council (NACC), the National Council for Population and Development (NCPD), and the Kenya Medical Research Institute (KEMRI). Funding for the KDHS was provided by the Government of Kenya with support from the United States Agency for International Development (USAID), the United Nations Population Fund (UNFPA), the United Kingdom Department for International Development (DfID), the World Bank, the Danish International Development Agency (DANIDA), the United Nations Children's Fund (UNICEF), the German Development Bank (KfW), the Clinton Health Access Initiative (CHAI), the World Food Programme (WFP), and the Micronutrient Initiative (MI). ICF International provided technical assistance as well as funding to the project through The DHS Program, a USAID-funded project providing support and technical assistance in the implementation of population and health surveys in countries worldwide.

Republic of Kenya

Additional information about the 2014 KDHS may be obtained from the Kenya National Bureau of Statistics (KNBS), P.O. Box 30266-00100 GPO Nairobi, Kenya; telephone (Nairobi): 3317586/8, 3317612/22, 3317623, 3317651; fax: 3315977; e-mail: directorgeneral@knbs.or.ke, info@knbs.or.ke; website: www.knbs.or.ke.

KNBS KENYA NATIONAL BUREAU OF STATISTICS *Keeping you informed*

Information on The DHS Program may be obtained from ICF International, 530 Gaither Road, Suite 500, Rockville MD, 20850, USA; telephone: 301-407-6500; fax: 301.407.6501; e-mail: info@DHSprogram.com; website: www.DHSprogram.com.



2014 Kenya DHS
Demographic and Health Survey

Murang'a


	Murang'a	Kenya total
Fertility and Family Planning		
Total fertility rate (number of children per woman)	3.0	3.9
Current use of *any* method of family planning (% of currently married women age 15-49)	70	58
Current use of any modern method of family planning (% of currently married women age 15-49)	63	53
Maternal and Child Health		
Maternity care		
Pregnant women who received antenatal care from a skilled provider[1] (%)	97	96
Pregnant women who received 4+ ANC visits (%)	61	58
Births assisted by a skilled provider[1] (%)	86	62
Births delivered in a health facility (%)	85	61
Child vaccination		
Children 12–23 months fully vaccinated[2] (%)	76	68
Nutrition		
Children under 5 years who are stunted (moderate or severe) (%)	19	26
Children under 5 years who are wasted (moderate or severe) (%)	1	4
Children under 5 years who are underweight (%)	6	11
Children under 5 years who are overweight or obese (%)	3	4
Malaria Prevention and Treatment		
Households with at least one insecticide-treated net[3] (ITN) (%)	44	59
Households with at least one ITN for every 2 persons in the household (universal coverage) (%)	27	34
Children under age 5 who slept under an ITN the night before the survey (%)	57	54
Pregnant women who slept under an ITN the night before the survey (%)	(45)	51
Pregnant women who received 3+ doses of SP/Fansidar to prevent malaria during pregnancy, at least one dose was received during an ANC visit (IPTp) (%)	2	10
Children under age 5 with fever in the 2 weeks before the survey	18	24
Among children under age 5 with fever, percentage who took any ACT	(0)	23
HIV/AIDS-related Knowledge		
Women 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	54	77
Men 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	78	85
Women 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	52	53
Men 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	46	46

Note: Figures in parentheses are based on 25-49 unweighted cases; an asterisk indicates that data are based on fewer than 25 unweighted cases and have been suppressed. [1]Skilled provider includes doctor, nurse, or midwife. [2]Fully vaccinated includes BCG, measles, three doses each of DPT and polio vaccine (excluding polio vaccine given at birth) and pneumococcal. [3]An insecticide-treated net (ITN) is (1) a factory-treated net that does not require any further treatment, or (2) a net that has been soaked with insecticide within the past 6 months.

Central and Nairobi: Key Indicators from the 2014 KDHS


Total Fertility Rate

Average number of live births per woman age 15-49




Use of Modern Methods of Family Planning

Percent of currently married women using a modern method of contraception




Antenatal Care: 4+ visits

Among women with a live birth in the 5 years before the survey, percent with 4+ ANC visits for their last live birth




Delivery in a Health Facility

Percentage of live births delivered in a health facility




Full Vaccination Rate

Percent of children age 12-23 months who have received all basic vaccinations plus the pneumococcal vaccine



ITN Access

Percentage of households with at least one insecticide-treated net (ITN) for every two persons who stayed in the household last night



The 2014 Kenya Demographic and Health Survey (2014 KDHS) was implemented by the Kenya National Bureau of Statistics from May 2014 to October 2014 in partnership with the Ministry of Health, the National AIDS Control Council (NACC), the National Council for Population and Development (NCPD), and the Kenya Medical Research Institute (KEMRI). Funding for the KDHS was provided by the Government of Kenya with support from the United States Agency for International Development (USAID), the United Nations Population Fund (UNFPA), the United Kingdom Department for International Development (DfID), the World Bank, the Danish International Development Agency (DANIDA), the United Nations Children's Fund (UNICEF), the German Development Bank (KfW), the Clinton Health Access Initiative (CHAI), the World Food Programme (WFP), and the Micronutrient Initiative (MI). ICF International provided technical assistance as well as funding to the project through The DHS Program, a USAID-funded project providing support and technical assistance in the implementation of population and health surveys in countries worldwide.

Republic of Kenya

Additional information about the 2014 KDHS may be obtained from the Kenya National Bureau of Statistics (KNBS), P.O. Box 30266-00100 GPO Nairobi, Kenya; telephone (Nairobi): 3317586/8, 3317612/22, 3317623, 3317651; fax: 3315977; e-mail: directorgeneral@knbs.or.ke, info@knbs.or.ke; website: www.knbs.or.ke.

KNBS KENYA NATIONAL BUREAU OF STATISTICS *Keeping you informed*

Information on The DHS Program may be obtained from ICF International, 530 Gaither Road, Suite 500, Rockville MD, 20850, USA; telephone: 301-407-6500; fax: 301.407.6501; e-mail: info@DHSprogram.com; website: www.DHSprogram.com.



2014 Kenya DHS
Demographic and Health Survey

Kiambu


	Kiambu	Kenya total
Fertility and Family Planning		
Total fertility rate (number of children per woman)	2.7	3.9
Current use of *any* method of family planning (% of currently married women age 15-49)	74	58
Current use of any modern method of family planning (% of currently married women age 15-49)	68	53
Maternal and Child Health		
Maternity care		
Pregnant women who received antenatal care from a skilled provider[1] (%)	98	96
Pregnant women who received 4+ ANC visits (%)	67	58
Births assisted by a skilled provider[1] (%)	93	62
Births delivered in a health facility (%)	93	61
Child vaccination		
Children 12–23 months fully vaccinated[2] (%)	83	68
Nutrition		
Children under 5 years who are stunted (moderate or severe) (%)	16	26
Children under 5 years who are wasted (moderate or severe) (%)	2	4
Children under 5 years who are underweight (%)	5	11
Children under 5 years who are overweight or obese (%)	8	4
Malaria Prevention and Treatment		
Households with at least one insecticide-treated net[3] (ITN) (%)	40	59
Households with at least one ITN for every 2 persons in the household (universal coverage) (%)	25	34
Children under age 5 who slept under an ITN the night before the survey (%)	48	54
Pregnant women who slept under an ITN the night before the survey (%)	(41)	51
Pregnant women who received 3+ doses of SP/Fansidar to prevent malaria during pregnancy, at least one dose was received during an ANC visit (IPTp) (%)	3	10
Children under age 5 with fever in the 2 weeks before the survey	19	24
Among children under age 5 with fever, percentage who took any ACT	1	23
HIV/AIDS-related Knowledge		
Women 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	76	77
Men 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	94	85
Women 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	52	53
Men 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	37	46

Note: Figures in parentheses are based on 25-49 unweighted cases; an asterisk indicates that data are based on fewer than 25 unweighted cases and have been suppressed. [1]Skilled provider includes doctor, nurse, or midwife. [2]Fully vaccinated includes BCG, measles, three doses each of DPT and polio vaccine (excluding polio vaccine given at birth) and pneumococcal. [3]An insecticide-treated net (ITN) is (1) a factory-treated net that does not require any further treatment, or (2) a net that has been soaked with insecticide within the past 6 months.

Central and Nairobi: Key Indicators from the 2014 KDHS


Total Fertility Rate

Average number of live births per woman age 15-49




Use of Modern Methods of Family Planning

Percent of currently married women using a modern method of contraception




Antenatal Care: 4+ visits

Among women with a live birth in the 5 years before the survey, percent with 4+ ANC visits for their last live birth




Delivery in a Health Facility

Percentage of live births delivered in a health facility




Full Vaccination Rate

Percent of children age 12-23 months who have received all basic vaccinations plus the pneumococcal vaccine



ITN Access

Percentage of households with at least one insecticide-treated net (ITN) for every two persons who stayed in the household last night




The 2014 Kenya Demographic and Health Survey (2014 KDHS) was implemented by the Kenya National Bureau of Statistics from May 2014 to October 2014 in partnership with the Ministry of Health, the National AIDS Control Council (NACC), the National Council for Population and Development (NCPD), and the Kenya Medical Research Institute (KEMRI). Funding for the KDHS was provided by the Government of Kenya with support from the United States Agency for International Development (USAID), the United Nations Population Fund (UNFPA), the United Kingdom Department for International Development (DfID), the World Bank, the Danish International Development Agency (DANIDA), the United Nations Children's Fund (UNICEF), the German Development Bank (KfW), the Clinton Health Access Initiative (CHAI), the World Food Programme (WFP), and the Micronutrient Initiative (MI). ICF International provided technical assistance as well as funding to the project through The DHS Program, a USAID-funded project providing support and technical assistance in the implementation of population and health surveys in countries worldwide.

Republic of Kenya

Additional information about the 2014 KDHS may be obtained from the Kenya National Bureau of Statistics (KNBS), P.O. Box 30266-00100 GPO Nairobi, Kenya; telephone (Nairobi): 3317586/8, 3317612/22, 3317623, 3317651; fax: 3315977; e-mail: directorgeneral@knbs.or.ke, info@knbs.or.ke; website: www.knbs.or.ke.

KNBS KENYA NATIONAL BUREAU OF STATISTICS *Keeping you informed*

Information on The DHS Program may be obtained from ICF International, 530 Gaither Road, Suite 500, Rockville MD, 20850, USA; telephone: 301-407-6500; fax: 301.407.6501; e-mail: info@DHSprogram.com; website: www.DHSprogram.com.



2014 Kenya DHS
Demographic and Health Survey

Turkana


	Turkana	Kenya total
Fertility and Family Planning		
Total fertility rate (number of children per woman)	6.9	3.9
Current use of *any* method of family planning (% of currently married women age 15-49)	10	58
Current use of any modern method of family planning (% of currently married women age 15-49)	10	53
Maternal and Child Health		
Maternity care		
Pregnant women who received antenatal care from a skilled provider[1] (%)	91	96
Pregnant women who received 4+ ANC visits (%)	49	58
Births assisted by a skilled provider[1] (%)	23	62
Births delivered in a health facility (%)	23	61
Child vaccination		
Children 12–23 months fully vaccinated[2] (%)	57	68
Nutrition		
Children under 5 years who are stunted (moderate or severe) (%)	24	26
Children under 5 years who are wasted (moderate or severe) (%)	23	4
Children under 5 years who are underweight (%)	34	11
Children under 5 years who are overweight or obese (%)	0.2	4
Malaria Prevention and Treatment		
Households with at least one insecticide-treated net[3] (ITN) (%)	46	59
Households with at least one ITN for every 2 persons in the household (universal coverage) (%)	16	34
Children under age 5 who slept under an ITN the night before the survey (%)	21	54
Pregnant women who slept under an ITN the night before the survey (%)	20	51
Pregnant women who received 3+ doses of SP/Fansidar to prevent malaria during pregnancy, at least one dose was received during an ANC visit (IPTp) (%)	13	10
Children under age 5 with fever in the 2 weeks before the survey	11	24
Among children under age 5 with fever, percentage who took any ACT	22	23
HIV/AIDS-related Knowledge		
Women 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	49	77
Men 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	2	85
Women 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	42	53
Men 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	60	46

Note: Figures in parentheses are based on 25-49 unweighted cases; an asterisk indicates that data are based on fewer than 25 unweighted cases and have been suppressed. [1]Skilled provider includes doctor, nurse, or midwife. [2]Fully vaccinated includes BCG, measles, three doses each of DPT and polio vaccine (excluding polio vaccine given at birth) and pneumococcal. [3]An insecticide-treated net (ITN) is (1) a factory-treated net that does not require any further treatment, or (2) a net that has been soaked with insecticide within the past 6 months.

Rift Valley: Key Indicators from the 2014 KDHS


Total Fertility Rate

Average number of live births per woman age 15-49



Use of Modern Methods of Family Planning


Percent of currently married women using a modern method of contraception



ANC and Place of Delivery

ANC: 4+ visits - Among women with a live birth in the 5 years before the survey, percent with 4+ ANC visits for their last live birth

Delivery in a health facility - Percentage of live births delivered in a health facility




The 2014 Kenya Demographic and Health Survey (2014 KDHS) was implemented by the Kenya National Bureau of Statistics from May 2014 to October 2014 in partnership with the Ministry of Health, the National AIDS Control Council (NACC), the National Council for Population and Development (NCPD), and the Kenya Medical Research Institute (KEMRI). Funding for the KDHS was provided by the Government of Kenya with support from the United States Agency for International Development (USAID), the United Nations Population Fund (UNFPA), the United Kingdom Department for International Development (DfID), the World Bank, the Danish International Development Agency (DANIDA), the United Nations Children's Fund (UNICEF), the German Development Bank (KfW), the Clinton Health Access Initiative (CHAI), the World Food Programme (WFP), and the Micronutrient Initiative (MI). ICF International provided technical assistance as well as funding to the project through The DHS Program, a USAID-funded project providing support and technical assistance in the implementation of population and health surveys in countries worldwide.

Republic of Kenya

Additional information about the 2014 KDHS may be obtained from the Kenya National Bureau of Statistics (KNBS), P.O. Box 30266-00100 GPO Nairobi, Kenya; telephone (Nairobi): 3317586/8, 3317612/22, 3317623, 3317651; fax: 3315977; e-mail: directorgeneral@knbs.or.ke, info@knbs.or.ke; website: www.knbs.or.ke.

Information on The DHS Program may be obtained from ICF International, 530 Gaither Road, Suite 500, Rockville MD, 20850, USA; telephone: 301-407-6500; fax: 301.407.6501; e-mail: info@DHSprogram.com; website: www.DHSprogram.com.



2014 Kenya DHS
Demographic and Health Survey

West Pokot


	West Pokot	Kenya total
Fertility and Family Planning		
Total fertility rate (number of children per woman)	7.2	3.9
Current use of *any* method of family planning (% of currently married women age 15-49)	14	58
Current use of any modern method of family planning (% of currently married women age 15-49)	13	53
Maternal and Child Health		
Maternity care		
Pregnant women who received antenatal care from a skilled provider[1] (%)	85	96
Pregnant women who received 4+ ANC visits (%)	18	58
Births assisted by a skilled provider[1] (%)	27	62
Births delivered in a health facility (%)	26	61
Child vaccination		
Children 12–23 months fully vaccinated[2] (%)	31	68
Nutrition		
Children under 5 years who are stunted (moderate or severe) (%)	46	26
Children under 5 years who are wasted (moderate or severe) (%)	14	4
Children under 5 years who are underweight (%)	39	11
Children under 5 years who are overweight or obese (%)	1	4
Malaria Prevention and Treatment		
Households with at least one insecticide-treated net[3] (ITN) (%)	60	59
Households with at least one ITN for every 2 persons in the household (universal coverage) (%)	21	34
Children under age 5 who slept under an ITN the night before the survey (%)	43	54
Pregnant women who slept under an ITN the night before the survey (%)	32	51
Pregnant women who received 3+ doses of SP/Fansidar to prevent malaria during pregnancy, at least one dose was received during an ANC visit (IPTp) (%)	3	10
Children under age 5 with fever in the 2 weeks before the survey	9	24
Among children under age 5 with fever, percentage who took any ACT	16	23
HIV/AIDS-related Knowledge		
Women 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	48	77
Men 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	77	85
Women 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	34	53
Men 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	20	46

Note: Figures in parentheses are based on 25-49 unweighted cases; an asterisk indicates that data are based on fewer than 25 unweighted cases and have been suppressed. [1]Skilled provider includes doctor, nurse, or midwife. [2]Fully vaccinated includes BCG, measles, three doses each of DPT and polio vaccine (excluding polio vaccine given at birth) and pneumococcal. [3]An insecticide-treated net (ITN) is (1) a factory-treated net that does not require any further treatment, or (2) a net that has been soaked with insecticide within the past 6 months.

Rift Valley: Key Indicators from the 2014 KDHS


Total Fertility Rate

Average number of live births per woman age 15-49



Use of Modern Methods of Family Planning


Percent of currently married women using a modern method of contraception



ANC and Place of Delivery

ANC: 4+ visits - Among women with a live birth in the 5 years before the survey, percent with 4+ ANC visits for their last live birth

Delivery in a health facility - Percentage of live births delivered in a health facility




The 2014 Kenya Demographic and Health Survey (2014 KDHS) was implemented by the Kenya National Bureau of Statistics from May 2014 to October 2014 in partnership with the Ministry of Health, the National AIDS Control Council (NACC), the National Council for Population and Development (NCPD), and the Kenya Medical Research Institute (KEMRI). Funding for the KDHS was provided by the Government of Kenya with support from the United States Agency for International Development (USAID), the United Nations Population Fund (UNFPA), the United Kingdom Department for International Development (DfID), the World Bank, the Danish International Development Agency (DANIDA), the United Nations Children's Fund (UNICEF), the German Development Bank (KfW), the Clinton Health Access Initiative (CHAI), the World Food Programme (WFP), and the Micronutrient Initiative (MI). ICF International provided technical assistance as well as funding to the project through The DHS Program, a USAID-funded project providing support and technical assistance in the implementation of population and health surveys in countries worldwide.

Republic of Kenya

Additional information about the 2014 KDHS may be obtained from the Kenya National Bureau of Statistics (KNBS), P.O. Box 30266-00100 GPO Nairobi, Kenya; telephone (Nairobi): 3317586/8, 3317612/22, 3317623, 3317651; fax: 3315977; e-mail: directorgeneral@knbs.or.ke, info@knbs.or.ke; website: www.knbs.or.ke.

Information on The DHS Program may be obtained from ICF International, 530 Gaither Road, Suite 500, Rockville MD, 20850, USA; telephone: 301-407-6500; fax: 301.407.6501; e-mail: info@DHSprogram.com; website: www.DHSprogram.com.



2014 Kenya DHS
Demographic and Health Survey

Samburu


	Samburu	Kenya total
Fertility and Family Planning		
Total fertility rate (number of children per woman)	6.3	3.9
Current use of *any* method of family planning (% of currently married women age 15-49)	23	58
Current use of any modern method of family planning (% of currently married women age 15-49)	20	53
Maternal and Child Health		
Maternity care		
Pregnant women who received antenatal care from a skilled provider[1] (%)	74	96
Pregnant women who received 4+ ANC visits (%)	52	58
Births assisted by a skilled provider[1] (%)	29	62
Births delivered in a health facility (%)	25	61
Child vaccination		
Children 12–23 months fully vaccinated[2] (%)	63	68
Nutrition		
Children under 5 years who are stunted (moderate or severe) (%)	30	26
Children under 5 years who are wasted (moderate or severe) (%)	14	4
Children under 5 years who are underweight (%)	29	11
Children under 5 years who are overweight or obese (%)	0.6	4
Malaria Prevention and Treatment		
Households with at least one insecticide-treated net[3] (ITN) (%)	19	59
Households with at least one ITN for every 2 persons in the household (universal coverage) (%)	9	34
Children under age 5 who slept under an ITN the night before the survey (%)	17	54
Pregnant women who slept under an ITN the night before the survey (%)	7	51
Pregnant women who received 3+ doses of SP/Fansidar to prevent malaria during pregnancy, at least one dose was received during an ANC visit (IPTp) (%)	3	10
Children under age 5 with fever in the 2 weeks before the survey	19	24
Among children under age 5 with fever, percentage who took any ACT	5	23
HIV/AIDS-related Knowledge		
Women 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	78	77
Men 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	79	85
Women 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	45	53
Men 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	51	46

Note: Figures in parentheses are based on 25-49 unweighted cases; an asterisk indicates that data are based on fewer than 25 unweighted cases and have been suppressed. [1]Skilled provider includes doctor, nurse, or midwife. [2]Fully vaccinated includes BCG, measles, three doses each of DPT and polio vaccine (excluding polio vaccine given at birth) and pneumococcal. [3]An insecticide-treated net (ITN) is (1) a factory-treated net that does not require any further treatment, or (2) a net that has been soaked with insecticide within the past 6 months.

Rift Valley: Key Indicators from the 2014 KDHS


Total Fertility Rate

Average number of live births per woman age 15-49



Use of Modern Methods of Family Planning


Percent of currently married women using a modern method of contraception



ANC and Place of Delivery

ANC: 4+ visits - Among women with a live birth in the 5 years before the survey, percent with 4+ ANC visits for their last live birth

Delivery in a health facility - Percentage of live births delivered in a health facility




The 2014 Kenya Demographic and Health Survey (2014 KDHS) was implemented by the Kenya National Bureau of Statistics from May 2014 to October 2014 in partnership with the Ministry of Health, the National AIDS Control Council (NACC), the National Council for Population and Development (NCPD), and the Kenya Medical Research Institute (KEMRI). Funding for the KDHS was provided by the Government of Kenya with support from the United States Agency for International Development (USAID), the United Nations Population Fund (UNFPA), the United Kingdom Department for International Development (DfID), the World Bank, the Danish International Development Agency (DANIDA), the United Nations Children's Fund (UNICEF), the German Development Bank (KfW), the Clinton Health Access Initiative (CHAI), the World Food Programme (WFP), and the Micronutrient Initiative (MI). ICF International provided technical assistance as well as funding to the project through The DHS Program, a USAID-funded project providing support and technical assistance in the implementation of population and health surveys in countries worldwide.

Republic of Kenya

Additional information about the 2014 KDHS may be obtained from the Kenya National Bureau of Statistics (KNBS), P.O. Box 30266-00100 GPO Nairobi, Kenya; telephone (Nairobi): 3317586/8, 3317612/22, 3317623, 3317651; fax: 3315977; e-mail: directorgeneral@knbs.or.ke, info@knbs.or.ke; website: www.knbs.or.ke.

Information on The DHS Program may be obtained from ICF International, 530 Gaither Road, Suite 500, Rockville MD, 20850, USA; telephone: 301-407-6500; fax: 301.407.6501; e-mail: info@DHSprogram.com; website: www.DHSprogram.com.



2014 Kenya DHS
Demographic and Health Survey

Trans-Nzoia


	Trans-Nzoia	Kenya total
Fertility and Family Planning		
Total fertility rate (number of children per woman)	5.2	3.9
Current use of *any* method of family planning (% of currently married women age 15-49)	64	58
Current use of any modern method of family planning (% of currently married women age 15-49)	56	53
Maternal and Child Health		
Maternity care		
Pregnant women who received antenatal care from a skilled provider[1] (%)	92	96
Pregnant women who received 4+ ANC visits (%)	41	58
Births assisted by a skilled provider[1] (%)	42	62
Births delivered in a health facility (%)	42	61
Child vaccination		
Children 12–23 months fully vaccinated[2] (%)	58	68
Nutrition		
Children under 5 years who are stunted (moderate or severe) (%)	29	26
Children under 5 years who are wasted (moderate or severe) (%)	4	4
Children under 5 years who are underweight (%)	15	11
Children under 5 years who are overweight or obese (%)	2	4
Malaria Prevention and Treatment		
Households with at least one insecticide-treated net[3] (ITN) (%)	71	59
Households with at least one ITN for every 2 persons in the household (universal coverage) (%)	38	34
Children under age 5 who slept under an ITN the night before the survey (%)	59	54
Pregnant women who slept under an ITN the night before the survey (%)	(56)	51
Pregnant women who received 3+ doses of SP/Fansidar to prevent malaria during pregnancy, at least one dose was received during an ANC visit (IPTp) (%)	3	10
Children under age 5 with fever in the 2 weeks before the survey	22	24
Among children under age 5 with fever, percentage who took any ACT	11	23
HIV/AIDS-related Knowledge		
Women 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	85	77
Men 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	80	85
Women 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	43	53
Men 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	41	46

Note: Figures in parentheses are based on 25-49 unweighted cases; an asterisk indicates that data are based on fewer than 25 unweighted cases and have been suppressed. [1]Skilled provider includes doctor, nurse, or midwife. [2]Fully vaccinated includes BCG, measles, three doses each of DPT and polio vaccine (excluding polio vaccine given at birth) and pneumococcal. [3]An insecticide-treated net (ITN) is (1) a factory-treated net that does not require any further treatment, or (2) a net that has been soaked with insecticide within the past 6 months.

Rift Valley: Key Indicators from the 2014 KDHS


Total Fertility Rate

Average number of live births per woman age 15-49



Use of Modern Methods of Family Planning


Percent of currently married women using a modern method of contraception



ANC and Place of Delivery

ANC: 4+ visits - Among women with a live birth in the 5 years before the survey, percent with 4+ ANC visits for their last live birth

Delivery in a health facility - Percentage of live births delivered in a health facility




The 2014 Kenya Demographic and Health Survey (2014 KDHS) was implemented by the Kenya National Bureau of Statistics from May 2014 to October 2014 in partnership with the Ministry of Health, the National AIDS Control Council (NACC), the National Council for Population and Development (NCPD), and the Kenya Medical Research Institute (KEMRI). Funding for the KDHS was provided by the Government of Kenya with support from the United States Agency for International Development (USAID), the United Nations Population Fund (UNFPA), the United Kingdom Department for International Development (DfID), the World Bank, the Danish International Development Agency (DANIDA), the United Nations Children's Fund (UNICEF), the German Development Bank (KfW), the Clinton Health Access Initiative (CHAI), the World Food Programme (WFP), and the Micronutrient Initiative (MI). ICF International provided technical assistance as well as funding to the project through The DHS Program, a USAID-funded project providing support and technical assistance in the implementation of population and health surveys in countries worldwide.

Republic of Kenya

Additional information about the 2014 KDHS may be obtained from the Kenya National Bureau of Statistics (KNBS), P.O. Box 30266-00100 GPO Nairobi, Kenya; telephone (Nairobi): 3317586/8, 3317612/22, 3317623, 3317651; fax: 3315977; e-mail: directorgeneral@knbs.or.ke, info@knbs.or.ke; website: www.knbs.or.ke.

Information on The DHS Program may be obtained from ICF International, 530 Gaither Road, Suite 500, Rockville MD, 20850, USA; telephone: 301-407-6500; fax: 301.407.6501; e-mail: info@DHSprogram.com; website: www.DHSprogram.com.



2014 Kenya DHS
Demographic and Health Survey

Uasin Gishu


	Uasin Gishu	Kenya total
Fertility and Family Planning		
Total fertility rate (number of children per woman)	3.6	3.9
Current use of *any* method of family planning (% of currently married women age 15-49)	63	58
Current use of any modern method of family planning (% of currently married women age 15-49)	56	53
Maternal and Child Health		
Maternity care		
Pregnant women who received antenatal care from a skilled provider[1] (%)	96	96
Pregnant women who received 4+ ANC visits (%)	63	58
Births assisted by a skilled provider[1] (%)	59	62
Births delivered in a health facility (%)	57	61
Child vaccination		
Children 12–23 months fully vaccinated[2] (%)	58	68
Nutrition		
Children under 5 years who are stunted (moderate or severe) (%)	31	26
Children under 5 years who are wasted (moderate or severe) (%)	3	4
Children under 5 years who are underweight (%)	12	11
Children under 5 years who are overweight or obese (%)	5	4
Malaria Prevention and Treatment		
Households with at least one insecticide-treated net[3] (ITN) (%)	72	59
Households with at least one ITN for every 2 persons in the household (universal coverage) (%)	50	34
Children under age 5 who slept under an ITN the night before the survey (%)	69	54
Pregnant women who slept under an ITN the night before the survey (%)	52	51
Pregnant women who received 3+ doses of SP/Fansidar to prevent malaria during pregnancy, at least one dose was received during an ANC visit (IPTp) (%)	6	10
Children under age 5 with fever in the 2 weeks before the survey	19	24
Among children under age 5 with fever, percentage who took any ACT	5	23
HIV/AIDS-related Knowledge		
Women 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	83	77
Men 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	83	85
Women 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	64	53
Men 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	54	46

Note: Figures in parentheses are based on 25-49 unweighted cases; an asterisk indicates that data are based on fewer than 25 unweighted cases and have been suppressed. [1]Skilled provider includes doctor, nurse, or midwife. [2]Fully vaccinated includes BCG, measles, three doses each of DPT and polio vaccine (excluding polio vaccine given at birth) and pneumococcal. [3]An insecticide-treated net (ITN) is (1) a factory-treated net that does not require any further treatment, or (2) a net that has been soaked with insecticide within the past 6 months.

Rift Valley: Key Indicators from the 2014 KDHS


Total Fertility Rate

Average number of live births per woman age 15-49



Use of Modern Methods of Family Planning


Percent of currently married women using a modern method of contraception



ANC and Place of Delivery

ANC: 4+ visits - Among women with a live birth in the 5 years before the survey, percent with 4+ ANC visits for their last live birth

Delivery in a health facility - Percentage of live births delivered in a health facility




The 2014 Kenya Demographic and Health Survey (2014 KDHS) was implemented by the Kenya National Bureau of Statistics from May 2014 to October 2014 in partnership with the Ministry of Health, the National AIDS Control Council (NACC), the National Council for Population and Development (NCPD), and the Kenya Medical Research Institute (KEMRI). Funding for the KDHS was provided by the Government of Kenya with support from the United States Agency for International Development (USAID), the United Nations Population Fund (UNFPA), the United Kingdom Department for International Development (DfID), the World Bank, the Danish International Development Agency (DANIDA), the United Nations Children's Fund (UNICEF), the German Development Bank (KfW), the Clinton Health Access Initiative (CHAI), the World Food Programme (WFP), and the Micronutrient Initiative (MI). ICF International provided technical assistance as well as funding to the project through The DHS Program, a USAID-funded project providing support and technical assistance in the implementation of population and health surveys in countries worldwide.

Republic of Kenya

Additional information about the 2014 KDHS may be obtained from the Kenya National Bureau of Statistics (KNBS), P.O. Box 30266-00100 GPO Nairobi, Kenya; telephone (Nairobi): 3317586/8, 3317612/22, 3317623, 3317651; fax: 3315977; e-mail: directorgeneral@knbs.or.ke, info@knbs.or.ke; website: www.knbs.or.ke.

Information on The DHS Program may be obtained from ICF International, 530 Gaither Road, Suite 500, Rockville MD, 20850, USA; telephone: 301-407-6500; fax: 301.407.6501; e-mail: info@DHSprogram.com; website: www.DHSprogram.com.



2014 Kenya DHS
Demographic and Health Survey

Elgeyo Marakwet


	Elgeyo Marakwet	Kenya total
Fertility and Family Planning		
Total fertility rate (number of children per woman)	4.1	3.9
Current use of *any* method of family planning (% of currently married women age 15-49)	55	58
Current use of any modern method of family planning (% of currently married women age 15-49)	44	53
Maternal and Child Health		
Maternity care		
Pregnant women who received antenatal care from a skilled provider[1] (%)	98	96
Pregnant women who received 4+ ANC visits (%)	47	58
Births assisted by a skilled provider[1] (%)	65	62
Births delivered in a health facility (%)	65	61
Child vaccination		
Children 12–23 months fully vaccinated[2] (%)	84	68
Nutrition		
Children under 5 years who are stunted (moderate or severe) (%)	30	26
Children under 5 years who are wasted (moderate or severe) (%)	4	4
Children under 5 years who are underweight (%)	13	11
Children under 5 years who are overweight or obese (%)	5	4
Malaria Prevention and Treatment		
Households with at least one insecticide-treated net[3] (ITN) (%)	22	59
Households with at least one ITN for every 2 persons in the household (universal coverage) (%)	10	34
Children under age 5 who slept under an ITN the night before the survey (%)	17	54
Pregnant women who slept under an ITN the night before the survey (%)	(26)	51
Pregnant women who received 3+ doses of SP/Fansidar to prevent malaria during pregnancy, at least one dose was received during an ANC visit (IPTp) (%)	2	10
Children under age 5 with fever in the 2 weeks before the survey	30	24
Among children under age 5 with fever, percentage who took any ACT	2	23
HIV/AIDS-related Knowledge		
Women 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	85	77
Men 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	97	85
Women 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	51	53
Men 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	34	46

Note: Figures in parentheses are based on 25-49 unweighted cases; an asterisk indicates that data are based on fewer than 25 unweighted cases and have been suppressed. [1]Skilled provider includes doctor, nurse, or midwife. [2]Fully vaccinated includes BCG, measles, three doses each of DPT and polio vaccine (excluding polio vaccine given at birth) and pneumococcal. [3]An insecticide-treated net (ITN) is (1) a factory-treated net that does not require any further treatment, or (2) a net that has been soaked with insecticide within the past 6 months.

Rift Valley: Key Indicators from the 2014 KDHS


Total Fertility Rate

Average number of live births per woman age 15-49



Use of Modern Methods of Family Planning


Percent of currently married women using a modern method of contraception



ANC and Place of Delivery

ANC: 4+ visits - Among women with a live birth in the 5 years before the survey, percent with 4+ ANC visits for their last live birth

Delivery in a health facility - Percentage of live births delivered in a health facility




The 2014 Kenya Demographic and Health Survey (2014 KDHS) was implemented by the Kenya National Bureau of Statistics from May 2014 to October 2014 in partnership with the Ministry of Health, the National AIDS Control Council (NACC), the National Council for Population and Development (NCPD), and the Kenya Medical Research Institute (KEMRI). Funding for the KDHS was provided by the Government of Kenya with support from the United States Agency for International Development (USAID), the United Nations Population Fund (UNFPA), the United Kingdom Department for International Development (DfID), the World Bank, the Danish International Development Agency (DANIDA), the United Nations Children's Fund (UNICEF), the German Development Bank (KfW), the Clinton Health Access Initiative (CHAI), the World Food Programme (WFP), and the Micronutrient Initiative (MI). ICF International provided technical assistance as well as funding to the project through The DHS Program, a USAID-funded project providing support and technical assistance in the implementation of population and health surveys in countries worldwide.

Republic of Kenya

Additional information about the 2014 KDHS may be obtained from the Kenya National Bureau of Statistics (KNBS), P.O. Box 30266-00100 GPO Nairobi, Kenya; telephone (Nairobi): 3317586/8, 3317612/22, 3317623, 3317651; fax: 3315977; e-mail: directorgeneral@knbs.or.ke, info@knbs.or.ke; website: www.knbs.or.ke.

Information on The DHS Program may be obtained from ICF International, 530 Gaither Road, Suite 500, Rockville MD, 20850, USA; telephone: 301-407-6500; fax: 301.407.6501; e-mail: info@DHSprogram.com; website: www.DHSprogram.com.



2014 Kenya DHS
Demographic and Health Survey

Nandi


	Nandi	Kenya total
Fertility and Family Planning		
Total fertility rate (number of children per woman)	4.0	3.9
Current use of *any* method of family planning (% of currently married women age 15-49)	65	58
Current use of any modern method of family planning (% of currently married women age 15-49)	59	53
Maternal and Child Health		
Maternity care		
Pregnant women who received antenatal care from a skilled provider[1] (%)	96	96
Pregnant women who received 4+ ANC visits (%)	57	58
Births assisted by a skilled provider[1] (%)	47	62
Births delivered in a health facility (%)	47	61
Child vaccination		
Children 12–23 months fully vaccinated[2] (%)	94	68
Nutrition		
Children under 5 years who are stunted (moderate or severe) (%)	30	26
Children under 5 years who are wasted (moderate or severe) (%)	4	4
Children under 5 years who are underweight (%)	11	11
Children under 5 years who are overweight or obese (%)	4	4
Malaria Prevention and Treatment		
Households with at least one insecticide-treated net[3] (ITN) (%)	79	59
Households with at least one ITN for every 2 persons in the household (universal coverage) (%)	42	34
Children under age 5 who slept under an ITN the night before the survey (%)	55	54
Pregnant women who slept under an ITN the night before the survey (%)	(62)	51
Pregnant women who received 3+ doses of SP/Fansidar to prevent malaria during pregnancy, at least one dose was received during an ANC visit (IPTp) (%)	2	10
Children under age 5 with fever in the 2 weeks before the survey	20	24
Among children under age 5 with fever, percentage who took any ACT	9	23
HIV/AIDS-related Knowledge		
Women 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	91	77
Men 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	98	85
Women 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	53	53
Men 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	54	46

Note: Figures in parentheses are based on 25-49 unweighted cases; an asterisk indicates that data are based on fewer than 25 unweighted cases and have been suppressed. [1]Skilled provider includes doctor, nurse, or midwife. [2]Fully vaccinated includes BCG, measles, three doses each of DPT and polio vaccine (excluding polio vaccine given at birth) and pneumococcal. [3]An insecticide-treated net (ITN) is (1) a factory-treated net that does not require any further treatment, or (2) a net that has been soaked with insecticide within the past 6 months.

Rift Valley: Key Indicators from the 2014 KDHS


Total Fertility Rate

Average number of live births per woman age 15-49



Use of Modern Methods of Family Planning


Percent of currently married women using a modern method of contraception



ANC and Place of Delivery

ANC: 4+ visits - Among women with a live birth in the 5 years before the survey, percent with 4+ ANC visits for their last live birth

Delivery in a health facility - Percentage of live births delivered in a health facility




The 2014 Kenya Demographic and Health Survey (2014 KDHS) was implemented by the Kenya National Bureau of Statistics from May 2014 to October 2014 in partnership with the Ministry of Health, the National AIDS Control Council (NACC), the National Council for Population and Development (NCPD), and the Kenya Medical Research Institute (KEMRI). Funding for the KDHS was provided by the Government of Kenya with support from the United States Agency for International Development (USAID), the United Nations Population Fund (UNFPA), the United Kingdom Department for International Development (DfID), the World Bank, the Danish International Development Agency (DANIDA), the United Nations Children's Fund (UNICEF), the German Development Bank (KfW), the Clinton Health Access Initiative (CHAI), the World Food Programme (WFP), and the Micronutrient Initiative (MI). ICF International provided technical assistance as well as funding to the project through The DHS Program, a USAID-funded project providing support and technical assistance in the implementation of population and health surveys in countries worldwide.

Republic of Kenya

Additional information about the 2014 KDHS may be obtained from the Kenya National Bureau of Statistics (KNBS), P.O. Box 30266-00100 GPO Nairobi, Kenya; telephone (Nairobi): 3317586/8, 3317612/22, 3317623, 3317651; fax: 3315977; e-mail: directorgeneral@knbs.or.ke, info@knbs.or.ke; website: www.knbs.or.ke.

Information on The DHS Program may be obtained from ICF International, 530 Gaither Road, Suite 500, Rockville MD, 20850, USA; telephone: 301-407-6500; fax: 301.407.6501; e-mail: info@DHSprogram.com; website: www.DHSprogram.com.



2014 Kenya DHS
Demographic and Health Survey

Baringo


	Baringo	Kenya total
Fertility and Family Planning		
Total fertility rate (number of children per woman)	4.8	3.9
Current use of *any* method of family planning (% of currently married women age 15-49)	41	58
Current use of any modern method of family planning (% of currently married women age 15-49)	33	53
Maternal and Child Health		
Maternity care		
Pregnant women who received antenatal care from a skilled provider[1] (%)	93	96
Pregnant women who received 4+ ANC visits (%)	55	58
Births assisted by a skilled provider[1] (%)	54	62
Births delivered in a health facility (%)	54	61
Child vaccination		
Children 12–23 months fully vaccinated[2] (%)	63	68
Nutrition		
Children under 5 years who are stunted (moderate or severe) (%)	30	26
Children under 5 years who are wasted (moderate or severe) (%)	7	4
Children under 5 years who are underweight (%)	20	11
Children under 5 years who are overweight or obese (%)	2	4
Malaria Prevention and Treatment		
Households with at least one insecticide-treated net[3] (ITN) (%)	60	59
Households with at least one ITN for every 2 persons in the household (universal coverage) (%)	33	34
Children under age 5 who slept under an ITN the night before the survey (%)	49	54
Pregnant women who slept under an ITN the night before the survey (%)	44	51
Pregnant women who received 3+ doses of SP/Fansidar to prevent malaria during pregnancy, at least one dose was received during an ANC visit (IPTp) (%)	7	10
Children under age 5 with fever in the 2 weeks before the survey	23	24
Among children under age 5 with fever, percentage who took any ACT	23	23
HIV/AIDS-related Knowledge		
Women 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	73	77
Men 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	90	85
Women 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	52	53
Men 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	44	46

Note: Figures in parentheses are based on 25-49 unweighted cases; an asterisk indicates that data are based on fewer than 25 unweighted cases and have been suppressed. [1]Skilled provider includes doctor, nurse, or midwife. [2]Fully vaccinated includes BCG, measles, three doses each of DPT and polio vaccine (excluding polio vaccine given at birth) and pneumococcal. [3]An insecticide-treated net (ITN) is (1) a factory-treated net that does not require any further treatment, or (2) a net that has been soaked with insecticide within the past 6 months.

Rift Valley: Key Indicators from the 2014 KDHS


Total Fertility Rate

Average number of live births per woman age 15-49



Use of Modern Methods of Family Planning


Percent of currently married women using a modern method of contraception



ANC and Place of Delivery

ANC: 4+ visits - Among women with a live birth in the 5 years before the survey, percent with 4+ ANC visits for their last live birth

Delivery in a health facility - Percentage of live births delivered in a health facility




The 2014 Kenya Demographic and Health Survey (2014 KDHS) was implemented by the Kenya National Bureau of Statistics from May 2014 to October 2014 in partnership with the Ministry of Health, the National AIDS Control Council (NACC), the National Council for Population and Development (NCPD), and the Kenya Medical Research Institute (KEMRI). Funding for the KDHS was provided by the Government of Kenya with support from the United States Agency for International Development (USAID), the United Nations Population Fund (UNFPA), the United Kingdom Department for International Development (DfID), the World Bank, the Danish International Development Agency (DANIDA), the United Nations Children's Fund (UNICEF), the German Development Bank (KfW), the Clinton Health Access Initiative (CHAI), the World Food Programme (WFP), and the Micronutrient Initiative (MI). ICF International provided technical assistance as well as funding to the project through The DHS Program, a USAID-funded project providing support and technical assistance in the implementation of population and health surveys in countries worldwide.

Republic of Kenya

Additional information about the 2014 KDHS may be obtained from the Kenya National Bureau of Statistics (KNBS), P.O. Box 30266-00100 GPO Nairobi, Kenya; telephone (Nairobi): 3317586/8, 3317612/22, 3317623, 3317651; fax: 3315977; e-mail: directorgeneral@knbs.or.ke, info@knbs.or.ke; website: www.knbs.or.ke.

Information on The DHS Program may be obtained from ICF International, 530 Gaither Road, Suite 500, Rockville MD, 20850, USA; telephone: 301-407-6500; fax: 301.407.6501; e-mail: info@DHSprogram.com; website: www.DHSprogram.com.



2014 Kenya DHS
Demographic and Health Survey

Laikipia


	Laikipia	Kenya total
Fertility and Family Planning		
Total fertility rate (number of children per woman)	3.7	3.9
Current use of *any* method of family planning (% of currently married women age 15-49)	59	58
Current use of any modern method of family planning (% of currently married women age 15-49)	51	53
Maternal and Child Health		
Maternity care		
Pregnant women who received antenatal care from a skilled provider[1] (%)	94	96
Pregnant women who received 4+ ANC visits (%)	56	58
Births assisted by a skilled provider[1] (%)	50	62
Births delivered in a health facility (%)	48	61
Child vaccination		
Children 12–23 months fully vaccinated[2] (%)	79	68
Nutrition		
Children under 5 years who are stunted (moderate or severe) (%)	27	26
Children under 5 years who are wasted (moderate or severe) (%)	4	4
Children under 5 years who are underweight (%)	14	11
Children under 5 years who are overweight or obese (%)	4	4
Malaria Prevention and Treatment		
Households with at least one insecticide-treated net[3] (ITN) (%)	18	59
Households with at least one ITN for every 2 persons in the household (universal coverage) (%)	11	34
Children under age 5 who slept under an ITN the night before the survey (%)	14	54
Pregnant women who slept under an ITN the night before the survey (%)	(7)	51
Pregnant women who received 3+ doses of SP/Fansidar to prevent malaria during pregnancy, at least one dose was received during an ANC visit (IPTp) (%)	2	10
Children under age 5 with fever in the 2 weeks before the survey	22	24
Among children under age 5 with fever, percentage who took any ACT	6	23
HIV/AIDS-related Knowledge		
Women 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	85	77
Men 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	72	85
Women 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	50	53
Men 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	43	46

Note: Figures in parentheses are based on 25-49 unweighted cases; an asterisk indicates that data are based on fewer than 25 unweighted cases and have been suppressed. [1]Skilled provider includes doctor, nurse, or midwife. [2]Fully vaccinated includes BCG, measles, three doses each of DPT and polio vaccine (excluding polio vaccine given at birth) and pneumococcal. [3]An insecticide-treated net (ITN) is (1) a factory-treated net that does not require any further treatment, or (2) a net that has been soaked with insecticide within the past 6 months.

Rift Valley: Key Indicators from the 2014 KDHS


Total Fertility Rate

Average number of live births per woman age 15-49



Use of Modern Methods of Family Planning


Percent of currently married women using a modern method of contraception



ANC and Place of Delivery

ANC: 4+ visits - Among women with a live birth in the 5 years before the survey, percent with 4+ ANC visits for their last live birth

Delivery in a health facility - Percentage of live births delivered in a health facility




The 2014 Kenya Demographic and Health Survey (2014 KDHS) was implemented by the Kenya National Bureau of Statistics from May 2014 to October 2014 in partnership with the Ministry of Health, the National AIDS Control Council (NACC), the National Council for Population and Development (NCPD), and the Kenya Medical Research Institute (KEMRI). Funding for the KDHS was provided by the Government of Kenya with support from the United States Agency for International Development (USAID), the United Nations Population Fund (UNFPA), the United Kingdom Department for International Development (DfID), the World Bank, the Danish International Development Agency (DANIDA), the United Nations Children's Fund (UNICEF), the German Development Bank (KfW), the Clinton Health Access Initiative (CHAI), the World Food Programme (WFP), and the Micronutrient Initiative (MI). ICF International provided technical assistance as well as funding to the project through The DHS Program, a USAID-funded project providing support and technical assistance in the implementation of population and health surveys in countries worldwide.

Republic of Kenya

Additional information about the 2014 KDHS may be obtained from the Kenya National Bureau of Statistics (KNBS), P.O. Box 30266-00100 GPO Nairobi, Kenya; telephone (Nairobi): 3317586/8, 3317612/22, 3317623, 3317651; fax: 3315977; e-mail: directorgeneral@knbs.or.ke, info@knbs.or.ke; website: www.knbs.or.ke.

Information on The DHS Program may be obtained from ICF International, 530 Gaither Road, Suite 500, Rockville MD, 20850, USA; telephone: 301-407-6500; fax: 301.407.6501; e-mail: info@DHSprogram.com; website: www.DHSprogram.com.



2014 Kenya DHS
Demographic and Health Survey

Nakuru


	Nakuru	Kenya total
Fertility and Family Planning		
Total fertility rate (number of children per woman)	3.7	3.9
Current use of *any* method of family planning (% of currently married women age 15-49)	57	58
Current use of any modern method of family planning (% of currently married women age 15-49)	54	53
Maternal and Child Health		
Maternity care		
Pregnant women who received antenatal care from a skilled provider[1] (%)	96	96
Pregnant women who received 4+ ANC visits (%)	61	58
Births assisted by a skilled provider[1] (%)	70	62
Births delivered in a health facility (%)	70	61
Child vaccination		
Children 12–23 months fully vaccinated[2] (%)	72	68
Nutrition		
Children under 5 years who are stunted (moderate or severe) (%)	28	26
Children under 5 years who are wasted (moderate or severe) (%)	5	4
Children under 5 years who are underweight (%)	10	11
Children under 5 years who are overweight or obese (%)	6	4
Malaria Prevention and Treatment		
Households with at least one insecticide-treated net[3] (ITN) (%)	38	59
Households with at least one ITN for every 2 persons in the household (universal coverage) (%)	23	34
Children under age 5 who slept under an ITN the night before the survey (%)	31	54
Pregnant women who slept under an ITN the night before the survey (%)	(35)	51
Pregnant women who received 3+ doses of SP/Fansidar to prevent malaria during pregnancy, at least one dose was received during an ANC visit (IPTp) (%)	4	10
Children under age 5 with fever in the 2 weeks before the survey	16	24
Among children under age 5 with fever, percentage who took any ACT	11	23
HIV/AIDS-related Knowledge		
Women 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	81	77
Men 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	91	85
Women 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	50	53
Men 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	49	46

Note: Figures in parentheses are based on 25-49 unweighted cases; an asterisk indicates that data are based on fewer than 25 unweighted cases and have been suppressed. [1]Skilled provider includes doctor, nurse, or midwife. [2]Fully vaccinated includes BCG, measles, three doses each of DPT and polio vaccine (excluding polio vaccine given at birth) and pneumococcal. [3]An insecticide-treated net (ITN) is (1) a factory-treated net that does not require any further treatment, or (2) a net that has been soaked with insecticide within the past 6 months.

Rift Valley: Key Indicators from the 2014 KDHS


Total Fertility Rate

Average number of live births per woman age 15-49



Use of Modern Methods of Family Planning


Percent of currently married women using a modern method of contraception



ANC and Place of Delivery

ANC: 4+ visits - Among women with a live birth in the 5 years before the survey, percent with 4+ ANC visits for their last live birth

Delivery in a health facility - Percentage of live births delivered in a health facility




The 2014 Kenya Demographic and Health Survey (2014 KDHS) was implemented by the Kenya National Bureau of Statistics from May 2014 to October 2014 in partnership with the Ministry of Health, the National AIDS Control Council (NACC), the National Council for Population and Development (NCPD), and the Kenya Medical Research Institute (KEMRI). Funding for the KDHS was provided by the Government of Kenya with support from the United States Agency for International Development (USAID), the United Nations Population Fund (UNFPA), the United Kingdom Department for International Development (DfID), the World Bank, the Danish International Development Agency (DANIDA), the United Nations Children's Fund (UNICEF), the German Development Bank (KfW), the Clinton Health Access Initiative (CHAI), the World Food Programme (WFP), and the Micronutrient Initiative (MI). ICF International provided technical assistance as well as funding to the project through The DHS Program, a USAID-funded project providing support and technical assistance in the implementation of population and health surveys in countries worldwide.

Republic of Kenya

Additional information about the 2014 KDHS may be obtained from the Kenya National Bureau of Statistics (KNBS), P.O. Box 30266-00100 GPO Nairobi, Kenya; telephone (Nairobi): 3317586/8, 3317612/22, 3317623, 3317651; fax: 3315977; e-mail: directorgeneral@knbs.or.ke, info@knbs.or.ke; website: www.knbs.or.ke.

Information on The DHS Program may be obtained from ICF International, 530 Gaither Road, Suite 500, Rockville MD, 20850, USA; telephone: 301-407-6500; fax: 301.407.6501; e-mail: info@DHSprogram.com; website: www.DHSprogram.com.



2014 Kenya DHS
Demographic and Health Survey

Narok


	Narok	Kenya total
Fertility and Family Planning		
Total fertility rate (number of children per woman)	6.0	3.9
Current use of *any* method of family planning (% of currently married women age 15-49)	48	58
Current use of any modern method of family planning (% of currently married women age 15-49)	38	53
Maternal and Child Health		
Maternity care		
Pregnant women who received antenatal care from a skilled provider[1] (%)	92	96
Pregnant women who received 4+ ANC visits (%)	46	58
Births assisted by a skilled provider[1] (%)	40	62
Births delivered in a health facility (%)	39	61
Child vaccination		
Children 12–23 months fully vaccinated[2] (%)	54	68
Nutrition		
Children under 5 years who are stunted (moderate or severe) (%)	33	26
Children under 5 years who are wasted (moderate or severe) (%)	2	4
Children under 5 years who are underweight (%)	12	11
Children under 5 years who are overweight or obese (%)	3	4
Malaria Prevention and Treatment		
Households with at least one insecticide-treated net[3] (ITN) (%)	52	59
Households with at least one ITN for every 2 persons in the household (universal coverage) (%)	25	34
Children under age 5 who slept under an ITN the night before the survey (%)	31	54
Pregnant women who slept under an ITN the night before the survey (%)	32	51
Pregnant women who received 3+ doses of SP/Fansidar to prevent malaria during pregnancy, at least one dose was received during an ANC visit (IPTp) (%)	0.6	10
Children under age 5 with fever in the 2 weeks before the survey	31	24
Among children under age 5 with fever, percentage who took any ACT	10	23
HIV/AIDS-related Knowledge		
Women 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	62	77
Men 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	78	85
Women 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	56	53
Men 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	43	46

Note: Figures in parentheses are based on 25-49 unweighted cases; an asterisk indicates that data are based on fewer than 25 unweighted cases and have been suppressed. [1]Skilled provider includes doctor, nurse, or midwife. [2]Fully vaccinated includes BCG, measles, three doses each of DPT and polio vaccine (excluding polio vaccine given at birth) and pneumococcal. [3]An insecticide-treated net (ITN) is (1) a factory-treated net that does not require any further treatment, or (2) a net that has been soaked with insecticide within the past 6 months.

Rift Valley: Key Indicators from the 2014 KDHS


Total Fertility Rate

Average number of live births per woman age 15-49



Use of Modern Methods of Family Planning


Percent of currently married women using a modern method of contraception



ANC and Place of Delivery

ANC: 4+ visits - Among women with a live birth in the 5 years before the survey, percent with 4+ ANC visits for their last live birth

Delivery in a health facility - Percentage of live births delivered in a health facility




The 2014 Kenya Demographic and Health Survey (2014 KDHS) was implemented by the Kenya National Bureau of Statistics from May 2014 to October 2014 in partnership with the Ministry of Health, the National AIDS Control Council (NACC), the National Council for Population and Development (NCPD), and the Kenya Medical Research Institute (KEMRI). Funding for the KDHS was provided by the Government of Kenya with support from the United States Agency for International Development (USAID), the United Nations Population Fund (UNFPA), the United Kingdom Department for International Development (DfID), the World Bank, the Danish International Development Agency (DANIDA), the United Nations Children's Fund (UNICEF), the German Development Bank (KfW), the Clinton Health Access Initiative (CHAI), the World Food Programme (WFP), and the Micronutrient Initiative (MI). ICF International provided technical assistance as well as funding to the project through The DHS Program, a USAID-funded project providing support and technical assistance in the implementation of population and health surveys in countries worldwide.

Republic of Kenya

Additional information about the 2014 KDHS may be obtained from the Kenya National Bureau of Statistics (KNBS), P.O. Box 30266-00100 GPO Nairobi, Kenya; telephone (Nairobi): 3317586/8, 3317612/22, 3317623, 3317651; fax: 3315977; e-mail: directorgeneral@knbs.or.ke, info@knbs.or.ke; website: www.knbs.or.ke.

Information on The DHS Program may be obtained from ICF International, 530 Gaither Road, Suite 500, Rockville MD, 20850, USA; telephone: 301-407-6500; fax: 301.407.6501; e-mail: info@DHSprogram.com; website: www.DHSprogram.com.



2014 Kenya DHS
Demographic and Health Survey

Kajiado


	Kajiado	Kenya total
Fertility and Family Planning		
Total fertility rate (number of children per woman)	4.5	3.9
Current use of *any* method of family planning (% of currently married women age 15-49)	55	58
Current use of any modern method of family planning (% of currently married women age 15-49)	45	53
Maternal and Child Health		
Maternity care		
Pregnant women who received antenatal care from a skilled provider[1] (%)	97	96
Pregnant women who received 4+ ANC visits (%)	67	58
Births assisted by a skilled provider[1] (%)	63	62
Births delivered in a health facility (%)	62	61
Child vaccination		
Children 12–23 months fully vaccinated[2] (%)	49	68
Nutrition		
Children under 5 years who are stunted (moderate or severe) (%)	18	26
Children under 5 years who are wasted (moderate or severe) (%)	3	4
Children under 5 years who are underweight (%)	8	11
Children under 5 years who are overweight or obese (%)	4	4
Malaria Prevention and Treatment		
Households with at least one insecticide-treated net[3] (ITN) (%)	50	59
Households with at least one ITN for every 2 persons in the household (universal coverage) (%)	33	34
Children under age 5 who slept under an ITN the night before the survey (%)	43	54
Pregnant women who slept under an ITN the night before the survey (%)	45	51
Pregnant women who received 3+ doses of SP/Fansidar to prevent malaria during pregnancy, at least one dose was received during an ANC visit (IPTp) (%)	9	10
Children under age 5 with fever in the 2 weeks before the survey	26	24
Among children under age 5 with fever, percentage who took any ACT	0	23
HIV/AIDS-related Knowledge		
Women 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	76	77
Men 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	85	85
Women 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	53	53
Men 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	53	46

Note: Figures in parentheses are based on 25-49 unweighted cases; an asterisk indicates that data are based on fewer than 25 unweighted cases and have been suppressed. [1]Skilled provider includes doctor, nurse, or midwife. [2]Fully vaccinated includes BCG, measles, three doses each of DPT and polio vaccine (excluding polio vaccine given at birth) and pneumococcal. [3]An insecticide-treated net (ITN) is (1) a factory-treated net that does not require any further treatment, or (2) a net that has been soaked with insecticide within the past 6 months.

Rift Valley: Key Indicators from the 2014 KDHS


Total Fertility Rate

Average number of live births per woman age 15-49



Use of Modern Methods of Family Planning


Percent of currently married women using a modern method of contraception



ANC and Place of Delivery

ANC: 4+ visits - Among women with a live birth in the 5 years before the survey, percent with 4+ ANC visits for their last live birth

Delivery in a health facility - Percentage of live births delivered in a health facility




The 2014 Kenya Demographic and Health Survey (2014 KDHS) was implemented by the Kenya National Bureau of Statistics from May 2014 to October 2014 in partnership with the Ministry of Health, the National AIDS Control Council (NACC), the National Council for Population and Development (NCPD), and the Kenya Medical Research Institute (KEMRI). Funding for the KDHS was provided by the Government of Kenya with support from the United States Agency for International Development (USAID), the United Nations Population Fund (UNFPA), the United Kingdom Department for International Development (DfID), the World Bank, the Danish International Development Agency (DANIDA), the United Nations Children's Fund (UNICEF), the German Development Bank (KfW), the Clinton Health Access Initiative (CHAI), the World Food Programme (WFP), and the Micronutrient Initiative (MI). ICF International provided technical assistance as well as funding to the project through The DHS Program, a USAID-funded project providing support and technical assistance in the implementation of population and health surveys in countries worldwide.

Republic of Kenya

Additional information about the 2014 KDHS may be obtained from the Kenya National Bureau of Statistics (KNBS), P.O. Box 30266-00100 GPO Nairobi, Kenya; telephone (Nairobi): 3317586/8, 3317612/22, 3317623, 3317651; fax: 3315977; e-mail: directorgeneral@knbs.or.ke, info@knbs.or.ke; website: www.knbs.or.ke.

Information on The DHS Program may be obtained from ICF International, 530 Gaither Road, Suite 500, Rockville MD, 20850, USA; telephone: 301-407-6500; fax: 301.407.6501; e-mail: info@DHSprogram.com; website: www.DHSprogram.com.



2014 Kenya DHS
Demographic and Health Survey

Kericho


	Kericho	Kenya total
Fertility and Family Planning		
Total fertility rate (number of children per woman)	4.0	3.9
Current use of *any* method of family planning (% of currently married women age 15-49)	63	58
Current use of any modern method of family planning (% of currently married women age 15-49)	57	53
Maternal and Child Health		
Maternity care		
Pregnant women who received antenatal care from a skilled provider[1] (%)	97	96
Pregnant women who received 4+ ANC visits (%)	54	58
Births assisted by a skilled provider[1] (%)	64	62
Births delivered in a health facility (%)	62	61
Child vaccination		
Children 12–23 months fully vaccinated[2] (%)	66	68
Nutrition		
Children under 5 years who are stunted (moderate or severe) (%)	29	26
Children under 5 years who are wasted (moderate or severe) (%)	6	4
Children under 5 years who are underweight (%)	12	11
Children under 5 years who are overweight or obese (%)	7	4
Malaria Prevention and Treatment		
Households with at least one insecticide-treated net[3] (ITN) (%)	80	59
Households with at least one ITN for every 2 persons in the household (universal coverage) (%)	45	34
Children under age 5 who slept under an ITN the night before the survey (%)	53	54
Pregnant women who slept under an ITN the night before the survey (%)	(68)	51
Pregnant women who received 3+ doses of SP/Fansidar to prevent malaria during pregnancy, at least one dose was received during an ANC visit (IPTp) (%)	3	10
Children under age 5 with fever in the 2 weeks before the survey	26	24
Among children under age 5 with fever, percentage who took any ACT	14	23
HIV/AIDS-related Knowledge		
Women 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	73	77
Men 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	90	85
Women 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	59	53
Men 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	51	46

Note: Figures in parentheses are based on 25-49 unweighted cases; an asterisk indicates that data are based on fewer than 25 unweighted cases and have been suppressed. [1]Skilled provider includes doctor, nurse, or midwife. [2]Fully vaccinated includes BCG, measles, three doses each of DPT and polio vaccine (excluding polio vaccine given at birth) and pneumococcal. [3]An insecticide-treated net (ITN) is (1) a factory-treated net that does not require any further treatment, or (2) a net that has been soaked with insecticide within the past 6 months.

Rift Valley: Key Indicators from the 2014 KDHS


Total Fertility Rate

Average number of live births per woman age 15-49



Use of Modern Methods of Family Planning


Percent of currently married women using a modern method of contraception



ANC and Place of Delivery

ANC: 4+ visits - Among women with a live birth in the 5 years before the survey, percent with 4+ ANC visits for their last live birth

Delivery in a health facility - Percentage of live births delivered in a health facility




The 2014 Kenya Demographic and Health Survey (2014 KDHS) was implemented by the Kenya National Bureau of Statistics from May 2014 to October 2014 in partnership with the Ministry of Health, the National AIDS Control Council (NACC), the National Council for Population and Development (NCPD), and the Kenya Medical Research Institute (KEMRI). Funding for the KDHS was provided by the Government of Kenya with support from the United States Agency for International Development (USAID), the United Nations Population Fund (UNFPA), the United Kingdom Department for International Development (DfID), the World Bank, the Danish International Development Agency (DANIDA), the United Nations Children's Fund (UNICEF), the German Development Bank (KfW), the Clinton Health Access Initiative (CHAI), the World Food Programme (WFP), and the Micronutrient Initiative (MI). ICF International provided technical assistance as well as funding to the project through The DHS Program, a USAID-funded project providing support and technical assistance in the implementation of population and health surveys in countries worldwide.

Republic of Kenya

Additional information about the 2014 KDHS may be obtained from the Kenya National Bureau of Statistics (KNBS), P.O. Box 30266-00100 GPO Nairobi, Kenya; telephone (Nairobi): 3317586/8, 3317612/22, 3317623, 3317651; fax: 3315977; e-mail: directorgeneral@knbs.or.ke, info@knbs.or.ke; website: www.knbs.or.ke.

Information on The DHS Program may be obtained from ICF International, 530 Gaither Road, Suite 500, Rockville MD, 20850, USA; telephone: 301-407-6500; fax: 301.407.6501; e-mail: info@DHSprogram.com; website: www.DHSprogram.com.



2014 Kenya DHS
Demographic and Health Survey

Bomet


	Bomet	Kenya total
Fertility and Family Planning		
Total fertility rate (number of children per woman)	4.3	3.9
Current use of *any* method of family planning (% of currently married women age 15-49)	55	58
Current use of any modern method of family planning (% of currently married women age 15-49)	50	53
Maternal and Child Health		
Maternity care		
Pregnant women who received antenatal care from a skilled provider[1] (%)	94	96
Pregnant women who received 4+ ANC visits (%)	37	58
Births assisted by a skilled provider[1] (%)	52	62
Births delivered in a health facility (%)	49	61
Child vaccination		
Children 12–23 months fully vaccinated[2] (%)	81	68
Nutrition		
Children under 5 years who are stunted (moderate or severe) (%)	36	26
Children under 5 years who are wasted (moderate or severe) (%)	2	4
Children under 5 years who are underweight (%)	12	11
Children under 5 years who are overweight or obese (%)	4	4
Malaria Prevention and Treatment		
Households with at least one insecticide-treated net[3] (ITN) (%)	78	59
Households with at least one ITN for every 2 persons in the household (universal coverage) (%)	47	34
Children under age 5 who slept under an ITN the night before the survey (%)	60	54
Pregnant women who slept under an ITN the night before the survey (%)	(48)	51
Pregnant women who received 3+ doses of SP/Fansidar to prevent malaria during pregnancy, at least one dose was received during an ANC visit (IPTp) (%)	5	10
Children under age 5 with fever in the 2 weeks before the survey	21	24
Among children under age 5 with fever, percentage who took any ACT	13	23
HIV/AIDS-related Knowledge		
Women 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	86	77
Men 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	90	85
Women 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	53	53
Men 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	44	46

Note: Figures in parentheses are based on 25-49 unweighted cases; an asterisk indicates that data are based on fewer than 25 unweighted cases and have been suppressed. [1]Skilled provider includes doctor, nurse, or midwife. [2]Fully vaccinated includes BCG, measles, three doses each of DPT and polio vaccine (excluding polio vaccine given at birth) and pneumococcal. [3]An insecticide-treated net (ITN) is (1) a factory-treated net that does not require any further treatment, or (2) a net that has been soaked with insecticide within the past 6 months.

Rift Valley: Key Indicators from the 2014 KDHS


Total Fertility Rate

Average number of live births per woman age 15-49



Use of Modern Methods of Family Planning


Percent of currently married women using a modern method of contraception



ANC and Place of Delivery

ANC: 4+ visits - Among women with a live birth in the 5 years before the survey, percent with 4+ ANC visits for their last live birth

Delivery in a health facility - Percentage of live births delivered in a health facility



The 2014 Kenya Demographic and Health Survey (2014 KDHS) was implemented by the Kenya National Bureau of Statistics from May 2014 to October 2014 in partnership with the Ministry of Health, the National AIDS Control Council (NACC), the National Council for Population and Development (NCPD), and the Kenya Medical Research Institute (KEMRI). Funding for the KDHS was provided by the Government of Kenya with support from the United States Agency for International Development (USAID), the United Nations Population Fund (UNFPA), the United Kingdom Department for International Development (DfID), the World Bank, the Danish International Development Agency (DANIDA), the United Nations Children's Fund (UNICEF), the German Development Bank (KfW), the Clinton Health Access Initiative (CHAI), the World Food Programme (WFP), and the Micronutrient Initiative (MI). ICF International provided technical assistance as well as funding to the project through The DHS Program, a USAID-funded project providing support and technical assistance in the implementation of population and health surveys in countries worldwide.

Republic of Kenya

Additional information about the 2014 KDHS may be obtained from the Kenya National Bureau of Statistics (KNBS), P.O. Box 30266-00100 GPO Nairobi, Kenya; telephone (Nairobi): 3317586/8, 3317612/22, 3317623, 3317651; fax: 3315977; e-mail: directorgeneral@knbs.or.ke, info@knbs.or.ke; website: www.knbs.or.ke.

Information on The DHS Program may be obtained from ICF International, 530 Gaither Road, Suite 500, Rockville MD, 20850, USA; telephone: 301-407-6500; fax: 301.407.6501; e-mail: info@DHSprogram.com; website: www.DHSprogram.com.



2014 Kenya DHS
Demographic and Health Survey

Kakamega


	Kakamega	Kenya total
Fertility and Family Planning		
Total fertility rate (number of children per woman)	4.4	3.9
Current use of *any* method of family planning (% of currently married women age 15-49)	62	58
Current use of any modern method of family planning (% of currently married women age 15-49)	60	53
Maternal and Child Health		
Maternity care		
Pregnant women who received antenatal care from a skilled provider[1] (%)	96	96
Pregnant women who received 4+ ANC visits (%)	45	58
Births assisted by a skilled provider[1] (%)	49	62
Births delivered in a health facility (%)	47	61
Child vaccination		
Children 12–23 months fully vaccinated[2] (%)	62	68
Nutrition		
Children under 5 years who are stunted (moderate or severe) (%)	28	26
Children under 5 years who are wasted (moderate or severe) (%)	2	4
Children under 5 years who are underweight (%)	10	11
Children under 5 years who are overweight or obese (%)	4	4
Malaria Prevention and Treatment		
Households with at least one insecticide-treated net[3] (ITN) (%)	79	59
Households with at least one ITN for every 2 persons in the household (universal coverage) (%)	47	34
Children under age 5 who slept under an ITN the night before the survey (%)	63	54
Pregnant women who slept under an ITN the night before the survey (%)	66	51
Pregnant women who received 3+ doses of SP/Fansidar to prevent malaria during pregnancy, at least one dose was received during an ANC visit (IPTp) (%)	20	10
Children under age 5 with fever in the 2 weeks before the survey	29	24
Among children under age 5 with fever, percentage who took any ACT	38	23
HIV/AIDS-related Knowledge		
Women 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	82	77
Men 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	83	85
Women 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	44	53
Men 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	42	46

Note: Figures in parentheses are based on 25-49 unweighted cases; an asterisk indicates that data are based on fewer than 25 unweighted cases and have been suppressed. [1]Skilled provider includes doctor, nurse, or midwife. [2]Fully vaccinated includes BCG, measles, three doses each of DPT and polio vaccine (excluding polio vaccine given at birth) and pneumococcal. [3]An insecticide-treated net (ITN) is (1) a factory-treated net that does not require any further treatment, or (2) a net that has been soaked with insecticide within the past 6 months.

Western: Key Indicators from the 2014 KDHS


Total Fertility Rate

Average number of live births per woman age 15-49




Use of Modern Methods of Family Planning

Percent of currently married women using a modern method of contraception




Antenatal Care: 4+ visits

Among women with a live birth in the 5 years before the survey, percent with 4+ ANC visits for their last live birth




Delivery in a Health Facility

Percentage of live births delivered in a health facility




Full Vaccination Rate

Percent of children age 12-23 months who have received all basic vaccinations plus the pneumococcal vaccine



ITN Access

Percentage of households with at least one insecticide-treated net (ITN) for every two persons who stayed in the household last night



The 2014 Kenya Demographic and Health Survey (2014 KDHS) was implemented by the Kenya National Bureau of Statistics from May 2014 to October 2014 in partnership with the Ministry of Health, the National AIDS Control Council (NACC), the National Council for Population and Development (NCPD), and the Kenya Medical Research Institute (KEMRI). Funding for the KDHS was provided by the Government of Kenya with support from the United States Agency for International Development (USAID), the United Nations Population Fund (UNFPA), the United Kingdom Department for International Development (DfID), the World Bank, the Danish International Development Agency (DANIDA), the United Nations Children's Fund (UNICEF), the German Development Bank (KfW), the Clinton Health Access Initiative (CHAI), the World Food Programme (WFP), and the Micronutrient Initiative (MI). ICF International provided technical assistance as well as funding to the project through The DHS Program, a USAID-funded project providing support and technical assistance in the implementation of population and health surveys in countries worldwide.

Republic of Kenya

Additional information about the 2014 KDHS may be obtained from the Kenya National Bureau of Statistics (KNBS), P.O. Box 30266-00100 GPO Nairobi, Kenya; telephone (Nairobi): 3317586/8, 3317612/22, 3317623, 3317651; fax: 3315977; e-mail: directorgeneral@knbs.or.ke, info@knbs.or.ke; website: www.knbs.or.ke.

Information on The DHS Program may be obtained from ICF International, 530 Gaither Road, Suite 500, Rockville MD, 20850, USA; telephone: 301-407-6500; fax: 301.407.6501; e-mail: info@DHSprogram.com; website: www.DHSprogram.com.



2014 KENYA DHS
Demographic and Health Survey

VIHIGA


	Vihiga	Kenya total
Fertility and Family Planning		
Total fertility rate (number of children per woman)	4.5	3.9
Current use of *any* method of family planning (% of currently married women age 15-49)	60	58
Current use of any modern method of family planning (% of currently married women age 15-49)	57	53
Maternal and Child Health		
Maternity care		
Pregnant women who received antenatal care from a skilled provider[1] (%)	97	96
Pregnant women who received 4+ ANC visits (%)	61	58
Births assisted by a skilled provider[1] (%)	50	62
Births delivered in a health facility (%)	50	61
Child vaccination		
Children 12–23 months fully vaccinated[2] (%)	91	68
Nutrition		
Children under 5 years who are stunted (moderate or severe) (%)	24	26
Children under 5 years who are wasted (moderate or severe) (%)	3	4
Children under 5 years who are underweight (%)	6	11
Children under 5 years who are overweight or obese (%)	4	4
Malaria Prevention and Treatment		
Households with at least one insecticide-treated net[3] (ITN) (%)	83	59
Households with at least one ITN for every 2 persons in the household (universal coverage) (%)	45	34
Children under age 5 who slept under an ITN the night before the survey (%)	71	54
Pregnant women who slept under an ITN the night before the survey (%)	(73)	51
Pregnant women who received 3+ doses of SP/Fansidar to prevent malaria during pregnancy, at least one dose was received during an ANC visit (IPTp) (%)	21	10
Children under age 5 with fever in the 2 weeks before the survey	49	24
Among children under age 5 with fever, percentage who took any ACT	40	23
HIV/AIDS-related Knowledge		
Women 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	78	77
Men 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	72	85
Women 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	45	53
Men 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	35	46

Note: Figures in parentheses are based on 25-49 unweighted cases; an asterisk indicates that data are based on fewer than 25 unweighted cases and have been suppressed. [1]Skilled provider includes doctor, nurse, or midwife. [2]Fully vaccinated includes BCG, measles, three doses each of DPT and polio vaccine (excluding polio vaccine given at birth) and pneumococcal. [3]An insecticide-treated net (ITN) is (1) a factory-treated net that does not require any further treatment, or (2) a net that has been soaked with insecticide within the past 6 months.

Western: Key Indicators from the 2014 KDHS


Total Fertility Rate

Average number of live births per woman age 15-49




Use of Modern Methods of Family Planning

Percent of currently married women using a modern method of contraception




Antenatal Care: 4+ visits

Among women with a live birth in the 5 years before the survey, percent with 4+ ANC visits for their last live birth




Delivery in a Health Facility

Percentage of live births delivered in a health facility




Full Vaccination Rate

Percent of children age 12-23 months who have received all basic vaccinations plus the pneumococcal vaccine



ITN Access

Percentage of households with at least one insecticide-treated net (ITN) for every two persons who stayed in the household last night



The 2014 Kenya Demographic and Health Survey (2014 KDHS) was implemented by the Kenya National Bureau of Statistics from May 2014 to October 2014 in partnership with the Ministry of Health, the National AIDS Control Council (NACC), the National Council for Population and Development (NCPD), and the Kenya Medical Research Institute (KEMRI). Funding for the KDHS was provided by the Government of Kenya with support from the United States Agency for International Development (USAID), the United Nations Population Fund (UNFPA), the United Kingdom Department for International Development (DfID), the World Bank, the Danish International Development Agency (DANIDA), the United Nations Children's Fund (UNICEF), the German Development Bank (KfW), the Clinton Health Access Initiative (CHAI), the World Food Programme (WFP), and the Micronutrient Initiative (MI). ICF International provided technical assistance as well as funding to the project through The DHS Program, a USAID-funded project providing support and technical assistance in the implementation of population and health surveys in countries worldwide.

Republic of Kenya

Additional information about the 2014 KDHS may be obtained from the Kenya National Bureau of Statistics (KNBS), P.O. Box 30266-00100 GPO Nairobi, Kenya; telephone (Nairobi): 3317586/8, 3317612/22, 3317623, 3317651; fax: 3315977; e-mail: directorgeneral@knbs.or.ke, info@knbs.or.ke; website: www.knbs.or.ke.

Information on The DHS Program may be obtained from ICF International, 530 Gaither Road, Suite 500, Rockville MD, 20850, USA; telephone: 301-407-6500; fax: 301.407.6501; e-mail: info@DHSprogram.com; website: www.DHSprogram.com.



2014 Kenya DHS
Demographic and Health Survey

Bungoma


	Bungoma	Kenya total
Fertility and Family Planning		
Total fertility rate (number of children per woman)	5.0	3.9
Current use of *any* method of family planning (% of currently married women age 15-49)	56	58
Current use of any modern method of family planning (% of currently married women age 15-49)	54	53
Maternal and Child Health		
Maternity care		
Pregnant women who received antenatal care from a skilled provider[1] (%)	98	96
Pregnant women who received 4+ ANC visits (%)	50	58
Births assisted by a skilled provider[1] (%)	41	62
Births delivered in a health facility (%)	41	61
Child vaccination		
Children 12–23 months fully vaccinated[2] (%)	73	68
Nutrition		
Children under 5 years who are stunted (moderate or severe) (%)	24	26
Children under 5 years who are wasted (moderate or severe) (%)	2	4
Children under 5 years who are underweight (%)	9	11
Children under 5 years who are overweight or obese (%)	3	4
Malaria Prevention and Treatment		
Households with at least one insecticide-treated net[3] (ITN) (%)	83	59
Households with at least one ITN for every 2 persons in the household (universal coverage) (%)	42	34
Children under age 5 who slept under an ITN the night before the survey (%)	72	54
Pregnant women who slept under an ITN the night before the survey (%)	62	51
Pregnant women who received 3+ doses of SP/Fansidar to prevent malaria during pregnancy, at least one dose was received during an ANC visit (IPTp) (%)	28	10
Children under age 5 with fever in the 2 weeks before the survey	36	24
Among children under age 5 with fever, percentage who took any ACT	55	23
HIV/AIDS-related Knowledge		
Women 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	84	77
Men 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	92	85
Women 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	44	53
Men 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	37	46

Note: Figures in parentheses are based on 25-49 unweighted cases; an asterisk indicates that data are based on fewer than 25 unweighted cases and have been suppressed. [1]Skilled provider includes doctor, nurse, or midwife. [2]Fully vaccinated includes BCG, measles, three doses each of DPT and polio vaccine (excluding polio vaccine given at birth) and pneumococcal. [3]An insecticide-treated net (ITN) is (1) a factory-treated net that does not require any further treatment, or (2) a net that has been soaked with insecticide within the past 6 months.

Western: Key Indicators from the 2014 KDHS


Total Fertility Rate

Average number of live births per woman age 15-49




Use of Modern Methods of Family Planning

Percent of currently married women using a modern method of contraception




Antenatal Care: 4+ visits

Among women with a live birth in the 5 years before the survey, percent with 4+ ANC visits for their last live birth




Delivery in a Health Facility

Percentage of live births delivered in a health facility




Full Vaccination Rate

Percent of children age 12-23 months who have received all basic vaccinations plus the pneumococcal vaccine



ITN Access

Percentage of households with at least one insecticide-treated net (ITN) for every two persons who stayed in the household last night



The 2014 Kenya Demographic and Health Survey (2014 KDHS) was implemented by the Kenya National Bureau of Statistics from May 2014 to October 2014 in partnership with the Ministry of Health, the National AIDS Control Council (NACC), the National Council for Population and Development (NCPD), and the Kenya Medical Research Institute (KEMRI). Funding for the KDHS was provided by the Government of Kenya with support from the United States Agency for International Development (USAID), the United Nations Population Fund (UNFPA), the United Kingdom Department for International Development (DfID), the World Bank, the Danish International Development Agency (DANIDA), the United Nations Children's Fund (UNICEF), the German Development Bank (KfW), the Clinton Health Access Initiative (CHAI), the World Food Programme (WFP), and the Micronutrient Initiative (MI). ICF International provided technical assistance as well as funding to the project through The DHS Program, a USAID-funded project providing support and technical assistance in the implementation of population and health surveys in countries worldwide.

Republic of Kenya

Additional information about the 2014 KDHS may be obtained from the Kenya National Bureau of Statistics (KNBS), P.O. Box 30266-00100 GPO Nairobi, Kenya; telephone (Nairobi): 3317586/8, 3317612/22, 3317623, 3317651; fax: 3315977; e-mail: directorgeneral@knbs.or.ke, info@knbs.or.ke; website: www.knbs.or.ke.

Information on The DHS Program may be obtained from ICF International, 530 Gaither Road, Suite 500, Rockville MD, 20850, USA; telephone: 301-407-6500; fax: 301.407.6501; e-mail: info@DHSprogram.com; website: www.DHSprogram.com.



2014 Kenya DHS
Demographic and Health Survey

Busia


	Busia	Kenya total
Fertility and Family Planning		
Total fertility rate (number of children per woman)	4.7	3.9
Current use of *any* method of family planning (% of currently married women age 15-49)	58	58
Current use of any modern method of family planning (% of currently married women age 15-49)	57	53
Maternal and Child Health		
Maternity care		
Pregnant women who received antenatal care from a skilled provider[1] (%)	98	96
Pregnant women who received 4+ ANC visits (%)	60	58
Births assisted by a skilled provider[1] (%)	59	62
Births delivered in a health facility (%)	58	61
Child vaccination		
Children 12–23 months fully vaccinated[2] (%)	75	68
Nutrition		
Children under 5 years who are stunted (moderate or severe) (%)	22	26
Children under 5 years who are wasted (moderate or severe) (%)	2	4
Children under 5 years who are underweight (%)	9	11
Children under 5 years who are overweight or obese (%)	2	4
Malaria Prevention and Treatment		
Households with at least one insecticide-treated net[3] (ITN) (%)	84	59
Households with at least one ITN for every 2 persons in the household (universal coverage) (%)	49	34
Children under age 5 who slept under an ITN the night before the survey (%)	74	54
Pregnant women who slept under an ITN the night before the survey (%)	(74)	51
Pregnant women who received 3+ doses of SP/Fansidar to prevent malaria during pregnancy, at least one dose was received during an ANC visit (IPTp) (%)	36	10
Children under age 5 with fever in the 2 weeks before the survey	43	24
Among children under age 5 with fever, percentage who took any ACT	60	23
HIV/AIDS-related Knowledge		
Women 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	76	77
Men 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	95	85
Women 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	50	53
Men 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	46	46

Note: Figures in parentheses are based on 25-49 unweighted cases; an asterisk indicates that data are based on fewer than 25 unweighted cases and have been suppressed. [1]Skilled provider includes doctor, nurse, or midwife. [2]Fully vaccinated includes BCG, measles, three doses each of DPT and polio vaccine (excluding polio vaccine given at birth) and pneumococcal. [3]An insecticide-treated net (ITN) is (1) a factory-treated net that does not require any further treatment, or (2) a net that has been soaked with insecticide within the past 6 months.

Western: Key Indicators from the 2014 KDHS


Total Fertility Rate

Average number of live births per woman age 15-49




Use of Modern Methods of Family Planning

Percent of currently married women using a modern method of contraception




Antenatal Care: 4+ visits

Among women with a live birth in the 5 years before the survey, percent with 4+ ANC visits for their last live birth




Delivery in a Health Facility

Percentage of live births delivered in a health facility




Full Vaccination Rate

Percent of children age 12-23 months who have received all basic vaccinations plus the pneumococcal vaccine



ITN Access

Percentage of households with at least one insecticide-treated net (ITN) for every two persons who stayed in the household last night



The 2014 Kenya Demographic and Health Survey (2014 KDHS) was implemented by the Kenya National Bureau of Statistics from May 2014 to October 2014 in partnership with the Ministry of Health, the National AIDS Control Council (NACC), the National Council for Population and Development (NCPD), and the Kenya Medical Research Institute (KEMRI). Funding for the KDHS was provided by the Government of Kenya with support from the United States Agency for International Development (USAID), the United Nations Population Fund (UNFPA), the United Kingdom Department for International Development (DfID), the World Bank, the Danish International Development Agency (DANIDA), the United Nations Children's Fund (UNICEF), the German Development Bank (KfW), the Clinton Health Access Initiative (CHAI), the World Food Programme (WFP), and the Micronutrient Initiative (MI). ICF International provided technical assistance as well as funding to the project through The DHS Program, a USAID-funded project providing support and technical assistance in the implementation of population and health surveys in countries worldwide.

Republic of Kenya

Additional information about the 2014 KDHS may be obtained from the Kenya National Bureau of Statistics (KNBS), P.O. Box 30266-00100 GPO Nairobi, Kenya; telephone (Nairobi): 3317586/8, 3317612/22, 3317623, 3317651; fax: 3315977; e-mail: directorgeneral@knbs.or.ke, info@knbs.or.ke; website: www.knbs.or.ke.

Information on The DHS Program may be obtained from ICF International, 530 Gaither Road, Suite 500, Rockville MD, 20850, USA; telephone: 301-407-6500; fax: 301.407.6501; e-mail: info@DHSprogram.com; website: www.DHSprogram.com.



2014 Kenya DHS
Demographic and Health Survey

Siaya


	Siaya	Kenya total
Fertility and Family Planning		
Total fertility rate (number of children per woman)	4.2	3.9
Current use of *any* method of family planning (% of currently married women age 15-49)	55	58
Current use of any modern method of family planning (% of currently married women age 15-49)	51	53
Maternal and Child Health		
Maternity care		
Pregnant women who received antenatal care from a skilled provider[1] (%)	98	96
Pregnant women who received 4+ ANC visits (%)	61	58
Births assisted by a skilled provider[1] (%)	70	62
Births delivered in a health facility (%)	70	61
Child vaccination		
Children 12–23 months fully vaccinated[2] (%)	73	68
Nutrition		
Children under 5 years who are stunted (moderate or severe) (%)	25	26
Children under 5 years who are wasted (moderate or severe) (%)	0.2	4
Children under 5 years who are underweight (%)	8	11
Children under 5 years who are overweight or obese (%)	5	4
Malaria Prevention and Treatment		
Households with at least one insecticide-treated net[3] (ITN) (%)	79	59
Households with at least one ITN for every 2 persons in the household (universal coverage) (%)	42	34
Children under age 5 who slept under an ITN the night before the survey (%)	68	54
Pregnant women who slept under an ITN the night before the survey (%)	(66)	51
Pregnant women who received 3+ doses of SP/Fansidar to prevent malaria during pregnancy, at least one dose was received during an ANC visit (IPTp) (%)	15	10
Children under age 5 with fever in the 2 weeks before the survey	45	24
Among children under age 5 with fever, percentage who took any ACT	59	23
HIV/AIDS-related Knowledge		
Women 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	87	77
Men 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	96	85
Women 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	58	53
Men 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	68	46

Note: Figures in parentheses are based on 25-49 unweighted cases; an asterisk indicates that data are based on fewer than 25 unweighted cases and have been suppressed. [1]Skilled provider includes doctor, nurse, or midwife. [2]Fully vaccinated includes BCG, measles, three doses each of DPT and polio vaccine (excluding polio vaccine given at birth) and pneumococcal. [3]An insecticide-treated net (ITN) is (1) a factory-treated net that does not require any further treatment, or (2) a net that has been soaked with insecticide within the past 6 months.

Nyanza: Key Indicators from the 2014 KDHS


Total Fertility Rate

Average number of live births per woman age 15-49




Use of Modern Methods of Family Planning

Percent of currently married women using a modern method of contraception




Antenatal Care: 4+ visits

Among women with a live birth in the 5 years before the survey, percent with 4+ ANC visits for their last live birth




Delivery in a Health Facility

Percentage of live births delivered in a health facility




Full Vaccination Rate

Percent of children age 12-23 months who have received all basic vaccinations plus the pneumococcal vaccine



ITN Access

Percentage of households with at least one insecticide-treated net (ITN) for every two persons who stayed in the household last night



The 2014 Kenya Demographic and Health Survey (2014 KDHS) was implemented by the Kenya National Bureau of Statistics from May 2014 to October 2014 in partnership with the Ministry of Health, the National AIDS Control Council (NACC), the National Council for Population and Development (NCPD), and the Kenya Medical Research Institute (KEMRI). Funding for the KDHS was provided by the Government of Kenya with support from the United States Agency for International Development (USAID), the United Nations Population Fund (UNFPA), the United Kingdom Department for International Development (DfID), the World Bank, the Danish International Development Agency (DANIDA), the United Nations Children's Fund (UNICEF), the German Development Bank (KfW), the Clinton Health Access Initiative (CHAI), the World Food Programme (WFP), and the Micronutrient Initiative (MI). ICF International provided technical assistance as well as funding to the project through The DHS Program, a USAID-funded project providing support and technical assistance in the implementation of population and health surveys in countries worldwide.

Republic of Kenya

Additional information about the 2014 KDHS may be obtained from the Kenya National Bureau of Statistics (KNBS), P.O. Box 30266-00100 GPO Nairobi, Kenya; telephone (Nairobi): 3317586/8, 3317612/22, 3317623, 3317651; fax: 3315977; e-mail: directorgeneral@knbs.or.ke, info@knbs.or.ke; website: www.knbs.or.ke.

KNBS KENYA NATIONAL BUREAU OF STATISTICS *Keeping you informed*

Information on The DHS Program may be obtained from ICF International, 530 Gaither Road, Suite 500, Rockville MD, 20850, USA; telephone: 301-407-6500; fax: 301.407.6501; e-mail: info@DHSprogram.com; website: www.DHSprogram.com.



2014 Kenya DHS
Demographic and Health Survey

Kisumu


	Kisumu	Kenya total
Fertility and Family Planning		
Total fertility rate (number of children per woman)	3.6	3.9
Current use of *any* method of family planning (% of currently married women age 15-49)	62	58
Current use of any modern method of family planning (% of currently married women age 15-49)	59	53
Maternal and Child Health		
Maternity care		
Pregnant women who received antenatal care from a skilled provider[1] (%)	98	96
Pregnant women who received 4+ ANC visits (%)	68	58
Births assisted by a skilled provider[1] (%)	69	62
Births delivered in a health facility (%)	70	61
Child vaccination		
Children 12–23 months fully vaccinated[2] (%)	73	68
Nutrition		
Children under 5 years who are stunted (moderate or severe) (%)	18	26
Children under 5 years who are wasted (moderate or severe) (%)	0.8	4
Children under 5 years who are underweight (%)	7	11
Children under 5 years who are overweight or obese (%)	6	4
Malaria Prevention and Treatment		
Households with at least one insecticide-treated net[3] (ITN) (%)	88	59
Households with at least one ITN for every 2 persons in the household (universal coverage) (%)	53	34
Children under age 5 who slept under an ITN the night before the survey (%)	82	54
Pregnant women who slept under an ITN the night before the survey (%)	(89)	51
Pregnant women who received 3+ doses of SP/Fansidar to prevent malaria during pregnancy, at least one dose was received during an ANC visit (IPTp) (%)	6	10
Children under age 5 with fever in the 2 weeks before the survey	31	24
Among children under age 5 with fever, percentage who took any ACT	40	23
HIV/AIDS-related Knowledge		
Women 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	84	77
Men 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	98	85
Women 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	62	53
Men 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	65	46

Note: Figures in parentheses are based on 25-49 unweighted cases; an asterisk indicates that data are based on fewer than 25 unweighted cases and have been suppressed. [1]Skilled provider includes doctor, nurse, or midwife. [2]Fully vaccinated includes BCG, measles, three doses each of DPT and polio vaccine (excluding polio vaccine given at birth) and pneumococcal. [3]An insecticide-treated net (ITN) is (1) a factory-treated net that does not require any further treatment, or (2) a net that has been soaked with insecticide within the past 6 months.

Nyanza: Key Indicators from the 2014 KDHS


Total Fertility Rate

Average number of live births per woman age 15-49




Use of Modern Methods of Family Planning

Percent of currently married women using a modern method of contraception




Antenatal Care: 4+ visits

Among women with a live birth in the 5 years before the survey, percent with 4+ ANC visits for their last live birth




Delivery in a Health Facility

Percentage of live births delivered in a health facility




Full Vaccination Rate

Percent of children age 12-23 months who have received all basic vaccinations plus the pneumococcal vaccine



ITN Access

Percentage of households with at least one insecticide-treated net (ITN) for every two persons who stayed in the household last night



The 2014 Kenya Demographic and Health Survey (2014 KDHS) was implemented by the Kenya National Bureau of Statistics from May 2014 to October 2014 in partnership with the Ministry of Health, the National AIDS Control Council (NACC), the National Council for Population and Development (NCPD), and the Kenya Medical Research Institute (KEMRI). Funding for the KDHS was provided by the Government of Kenya with support from the United States Agency for International Development (USAID), the United Nations Population Fund (UNFPA), the United Kingdom Department for International Development (DfID), the World Bank, the Danish International Development Agency (DANIDA), the United Nations Children's Fund (UNICEF), the German Development Bank (KfW), the Clinton Health Access Initiative (CHAI), the World Food Programme (WFP), and the Micronutrient Initiative (MI). ICF International provided technical assistance as well as funding to the project through The DHS Program, a USAID-funded project providing support and technical assistance in the implementation of population and health surveys in countries worldwide.

Republic of Kenya

Additional information about the 2014 KDHS may be obtained from the Kenya National Bureau of Statistics (KNBS), P.O. Box 30266-00100 GPO Nairobi, Kenya; telephone (Nairobi): 3317586/8, 3317612/22, 3317623, 3317651; fax: 3315977; e-mail: directorgeneral@knbs.or.ke, info@knbs.or.ke; website: www.knbs.or.ke.

KNBS KENYA NATIONAL BUREAU OF STATISTICS *Keeping you informed*

Information on The DHS Program may be obtained from ICF International, 530 Gaither Road, Suite 500, Rockville MD, 20850, USA; telephone: 301-407-6500; fax: 301.407.6501; e-mail: info@DHSprogram.com; website: www.DHSprogram.com.



2014 Kenya DHS
Demographic and Health Survey

Homa Bay


	Homa Bay	Kenya total
Fertility and Family Planning		
Total fertility rate (number of children per woman)	5.2	3.9
Current use of *any* method of family planning (% of currently married women age 15-49)	47	58
Current use of any modern method of family planning (% of currently married women age 15-49)	46	53
Maternal and Child Health		
Maternity care		
Pregnant women who received antenatal care from a skilled provider[1] (%)	94	96
Pregnant women who received 4+ ANC visits (%)	61	58
Births assisted by a skilled provider[1] (%)	60	62
Births delivered in a health facility (%)	62	61
Child vaccination		
Children 12–23 months fully vaccinated[2] (%)	54	68
Nutrition		
Children under 5 years who are stunted (moderate or severe) (%)	19	26
Children under 5 years who are wasted (moderate or severe) (%)	2	4
Children under 5 years who are underweight (%)	5	11
Children under 5 years who are overweight or obese (%)	4	4
Malaria Prevention and Treatment		
Households with at least one insecticide-treated net[3] (ITN) (%)	74	59
Households with at least one ITN for every 2 persons in the household (universal coverage) (%)	38	34
Children under age 5 who slept under an ITN the night before the survey (%)	56	54
Pregnant women who slept under an ITN the night before the survey (%)	(66)	51
Pregnant women who received 3+ doses of SP/Fansidar to prevent malaria during pregnancy, at least one dose was received during an ANC visit (IPTp) (%)	12	10
Children under age 5 with fever in the 2 weeks before the survey	46	24
Among children under age 5 with fever, percentage who took any ACT	47	23
HIV/AIDS-related Knowledge		
Women 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	88	77
Men 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	97	85
Women 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	71	53
Men 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	59	46

Note: Figures in parentheses are based on 25-49 unweighted cases; an asterisk indicates that data are based on fewer than 25 unweighted cases and have been suppressed. [1]Skilled provider includes doctor, nurse, or midwife. [2]Fully vaccinated includes BCG, measles, three doses each of DPT and polio vaccine (excluding polio vaccine given at birth) and pneumococcal. [3]An insecticide-treated net (ITN) is (1) a factory-treated net that does not require any further treatment, or (2) a net that has been soaked with insecticide within the past 6 months.

Nyanza: Key Indicators from the 2014 KDHS


Total Fertility Rate

Average number of live births per woman age 15-49




Use of Modern Methods of Family Planning

Percent of currently married women using a modern method of contraception




Antenatal Care: 4+ visits

Among women with a live birth in the 5 years before the survey, percent with 4+ ANC visits for their last live birth




Delivery in a Health Facility

Percentage of live births delivered in a health facility




Full Vaccination Rate

Percent of children age 12-23 months who have received all basic vaccinations plus the pneumococcal vaccine



ITN Access

Percentage of households with at least one insecticide-treated net (ITN) for every two persons who stayed in the household last night



The 2014 Kenya Demographic and Health Survey (2014 KDHS) was implemented by the Kenya National Bureau of Statistics from May 2014 to October 2014 in partnership with the Ministry of Health, the National AIDS Control Council (NACC), the National Council for Population and Development (NCPD), and the Kenya Medical Research Institute (KEMRI). Funding for the KDHS was provided by the Government of Kenya with support from the United States Agency for International Development (USAID), the United Nations Population Fund (UNFPA), the United Kingdom Department for International Development (DfID), the World Bank, the Danish International Development Agency (DANIDA), the United Nations Children's Fund (UNICEF), the German Development Bank (KfW), the Clinton Health Access Initiative (CHAI), the World Food Programme (WFP), and the Micronutrient Initiative (MI). ICF International provided technical assistance as well as funding to the project through The DHS Program, a USAID-funded project providing support and technical assistance in the implementation of population and health surveys in countries worldwide.

Republic of Kenya

Additional information about the 2014 KDHS may be obtained from the Kenya National Bureau of Statistics (KNBS), P.O. Box 30266-00100 GPO Nairobi, Kenya; telephone (Nairobi): 3317586/8, 3317612/22, 3317623, 3317651; fax: 3315977; e-mail: directorgeneral@knbs.or.ke, info@knbs.or.ke; website: www.knbs.or.ke.

KNBS KENYA NATIONAL BUREAU OF STATISTICS *Keeping you informed*

Information on The DHS Program may be obtained from ICF International, 530 Gaither Road, Suite 500, Rockville MD, 20850, USA; telephone: 301-407-6500; fax: 301.407.6501; e-mail: info@DHSprogram.com; website: www.DHSprogram.com.



2014 Kenya DHS
Demographic and Health Survey

Migori


	Migori	Kenya total
Fertility and Family Planning		
Total fertility rate (number of children per woman)	5.3	3.9
Current use of *any* method of family planning (% of currently married women age 15-49)	45	58
Current use of any modern method of family planning (% of currently married women age 15-49)	44	53
Maternal and Child Health		
Maternity care		
Pregnant women who received antenatal care from a skilled provider[1] (%)	96	96
Pregnant women who received 4+ ANC visits (%)	56	58
Births assisted by a skilled provider[1] (%)	53	62
Births delivered in a health facility (%)	53	61
Child vaccination		
Children 12–23 months fully vaccinated[2] (%)	37	68
Nutrition		
Children under 5 years who are stunted (moderate or severe) (%)	26	26
Children under 5 years who are wasted (moderate or severe) (%)	4	4
Children under 5 years who are underweight (%)	9	11
Children under 5 years who are overweight or obese (%)	4	4
Malaria Prevention and Treatment		
Households with at least one insecticide-treated net[3] (ITN) (%)	75	59
Households with at least one ITN for every 2 persons in the household (universal coverage) (%)	30	34
Children under age 5 who slept under an ITN the night before the survey (%)	62	54
Pregnant women who slept under an ITN the night before the survey (%)	56	51
Pregnant women who received 3+ doses of SP/Fansidar to prevent malaria during pregnancy, at least one dose was received during an ANC visit (IPTp) (%)	13	10
Children under age 5 with fever in the 2 weeks before the survey	48	24
Among children under age 5 with fever, percentage who took any ACT	37	23
HIV/AIDS-related Knowledge		
Women 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	80	77
Men 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	80	85
Women 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	65	53
Men 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	62	46

Note: Figures in parentheses are based on 25-49 unweighted cases; an asterisk indicates that data are based on fewer than 25 unweighted cases and have been suppressed. [1]Skilled provider includes doctor, nurse, or midwife. [2]Fully vaccinated includes BCG, measles, three doses each of DPT and polio vaccine (excluding polio vaccine given at birth) and pneumococcal. [3]An insecticide-treated net (ITN) is (1) a factory-treated net that does not require any further treatment, or (2) a net that has been soaked with insecticide within the past 6 months.

Nyanza: Key Indicators from the 2014 KDHS


Total Fertility Rate

Average number of live births per woman age 15-49




Use of Modern Methods of Family Planning

Percent of currently married women using a modern method of contraception




Antenatal Care: 4+ visits

Among women with a live birth in the 5 years before the survey, percent with 4+ ANC visits for their last live birth




Delivery in a Health Facility

Percentage of live births delivered in a health facility




Full Vaccination Rate

Percent of children age 12-23 months who have received all basic vaccinations plus the pneumococcal vaccine



ITN Access

Percentage of households with at least one insecticide-treated net (ITN) for every two persons who stayed in the household last night



The 2014 Kenya Demographic and Health Survey (2014 KDHS) was implemented by the Kenya National Bureau of Statistics from May 2014 to October 2014 in partnership with the Ministry of Health, the National AIDS Control Council (NACC), the National Council for Population and Development (NCPD), and the Kenya Medical Research Institute (KEMRI). Funding for the KDHS was provided by the Government of Kenya with support from the United States Agency for International Development (USAID), the United Nations Population Fund (UNFPA), the United Kingdom Department for International Development (DfID), the World Bank, the Danish International Development Agency (DANIDA), the United Nations Children's Fund (UNICEF), the German Development Bank (KfW), the Clinton Health Access Initiative (CHAI), the World Food Programme (WFP), and the Micronutrient Initiative (MI). ICF International provided technical assistance as well as funding to the project through The DHS Program, a USAID-funded project providing support and technical assistance in the implementation of population and health surveys in countries worldwide.

Republic of Kenya

Additional information about the 2014 KDHS may be obtained from the Kenya National Bureau of Statistics (KNBS), P.O. Box 30266-00100 GPO Nairobi, Kenya; telephone (Nairobi): 3317586/8, 3317612/22, 3317623, 3317651; fax: 3315977; e-mail: directorgeneral@knbs.or.ke, info@knbs.or.ke; website: www.knbs.or.ke.

KNBS KENYA NATIONAL BUREAU OF STATISTICS *Keeping you informed*

Information on The DHS Program may be obtained from ICF International, 530 Gaither Road, Suite 500, Rockville MD, 20850, USA; telephone: 301-407-6500; fax: 301.407.6501; e-mail: info@DHSprogram.com; website: www.DHSprogram.com.



2014 Kenya DHS
Demographic and Health Survey

Kisii


	Kisii	Kenya total
Fertility and Family Planning		
Total fertility rate (number of children per woman)	3.7	3.9
Current use of *any* method of family planning (% of currently married women age 15-49)	66	58
Current use of any modern method of family planning (% of currently married women age 15-49)	63	53
Maternal and Child Health		
Maternity care		
Pregnant women who received antenatal care from a skilled provider[1] (%)	98	96
Pregnant women who received 4+ ANC visits (%)	50	58
Births assisted by a skilled provider[1] (%)	73	62
Births delivered in a health facility (%)	69	61
Child vaccination		
Children 12–23 months fully vaccinated[2] (%)	78	68
Nutrition		
Children under 5 years who are stunted (moderate or severe) (%)	26	26
Children under 5 years who are wasted (moderate or severe) (%)	2	4
Children under 5 years who are underweight (%)	9	11
Children under 5 years who are overweight or obese (%)	4	4
Malaria Prevention and Treatment		
Households with at least one insecticide-treated net[3] (ITN) (%)	86	59
Households with at least one ITN for every 2 persons in the household (universal coverage) (%)	54	34
Children under age 5 who slept under an ITN the night before the survey (%)	80	54
Pregnant women who slept under an ITN the night before the survey (%)	(77)	51
Pregnant women who received 3+ doses of SP/Fansidar to prevent malaria during pregnancy, at least one dose was received during an ANC visit (IPTp) (%)	4	10
Children under age 5 with fever in the 2 weeks before the survey	28	24
Among children under age 5 with fever, percentage who took any ACT	27	23
HIV/AIDS-related Knowledge		
Women 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	80	77
Men 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	89	85
Women 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	51	53
Men 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	39	46

Note: Figures in parentheses are based on 25-49 unweighted cases; an asterisk indicates that data are based on fewer than 25 unweighted cases and have been suppressed. [1]Skilled provider includes doctor, nurse, or midwife. [2]Fully vaccinated includes BCG, measles, three doses each of DPT and polio vaccine (excluding polio vaccine given at birth) and pneumococcal. [3]An insecticide-treated net (ITN) is (1) a factory-treated net that does not require any further treatment, or (2) a net that has been soaked with insecticide within the past 6 months.

Nyanza: Key Indicators from the 2014 KDHS


Total Fertility Rate

Average number of live births per woman age 15-49




Use of Modern Methods of Family Planning

Percent of currently married women using a modern method of contraception




Antenatal Care: 4+ visits

Among women with a live birth in the 5 years before the survey, percent with 4+ ANC visits for their last live birth




Delivery in a Health Facility

Percentage of live births delivered in a health facility




Full Vaccination Rate

Percent of children age 12-23 months who have received all basic vaccinations plus the pneumococcal vaccine



ITN Access

Percentage of households with at least one insecticide-treated net (ITN) for every two persons who stayed in the household last night



The 2014 Kenya Demographic and Health Survey (2014 KDHS) was implemented by the Kenya National Bureau of Statistics from May 2014 to October 2014 in partnership with the Ministry of Health, the National AIDS Control Council (NACC), the National Council for Population and Development (NCPD), and the Kenya Medical Research Institute (KEMRI). Funding for the KDHS was provided by the Government of Kenya with support from the United States Agency for International Development (USAID), the United Nations Population Fund (UNFPA), the United Kingdom Department for International Development (DfID), the World Bank, the Danish International Development Agency (DANIDA), the United Nations Children's Fund (UNICEF), the German Development Bank (KfW), the Clinton Health Access Initiative (CHAI), the World Food Programme (WFP), and the Micronutrient Initiative (MI). ICF International provided technical assistance as well as funding to the project through The DHS Program, a USAID-funded project providing support and technical assistance in the implementation of population and health surveys in countries worldwide.

Republic of Kenya

Additional information about the 2014 KDHS may be obtained from the Kenya National Bureau of Statistics (KNBS), P.O. Box 30266-00100 GPO Nairobi, Kenya; telephone (Nairobi): 3317586/8, 3317612/22, 3317623, 3317651; fax: 3315977; e-mail: directorgeneral@knbs.or.ke, info@knbs.or.ke; website: www.knbs.or.ke.

KNBS KENYA NATIONAL BUREAU OF STATISTICS *Keeping you informed*

Information on The DHS Program may be obtained from ICF International, 530 Gaither Road, Suite 500, Rockville MD, 20850, USA; telephone: 301-407-6500; fax: 301.407.6501; e-mail: info@DHSprogram.com; website: www.DHSprogram.com.



2014 Kenya DHS
Demographic and Health Survey

Nyamira


	Nyamira	Kenya total
Fertility and Family Planning		
Total fertility rate (number of children per woman)	3.5	3.9
Current use of *any* method of family planning (% of currently married women age 15-49)	68	58
Current use of any modern method of family planning (% of currently married women age 15-49)	64	53
Maternal and Child Health		
Maternity care		
Pregnant women who received antenatal care from a skilled provider[1] (%)	96	96
Pregnant women who received 4+ ANC visits (%)	50	58
Births assisted by a skilled provider[1] (%)	74	62
Births delivered in a health facility (%)	74	61
Child vaccination		
Children 12–23 months fully vaccinated[2] (%)	87	68
Nutrition		
Children under 5 years who are stunted (moderate or severe) (%)	26	26
Children under 5 years who are wasted (moderate or severe) (%)	4	4
Children under 5 years who are underweight (%)	10	11
Children under 5 years who are overweight or obese (%)	3	4
Malaria Prevention and Treatment		
Households with at least one insecticide-treated net[3] (ITN) (%)	85	59
Households with at least one ITN for every 2 persons in the household (universal coverage) (%)	57	34
Children under age 5 who slept under an ITN the night before the survey (%)	70	54
Pregnant women who slept under an ITN the night before the survey (%)	*	51
Pregnant women who received 3+ doses of SP/Fansidar to prevent malaria during pregnancy, at least one dose was received during an ANC visit (IPTp) (%)	7	10
Children under age 5 with fever in the 2 weeks before the survey	6	24
Among children under age 5 with fever, percentage who took any ACT	*	23
HIV/AIDS-related Knowledge		
Women 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	98	77
Men 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	98	85
Women 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	51	53
Men 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	42	46

Note: Figures in parentheses are based on 25-49 unweighted cases; an asterisk indicates that data are based on fewer than 25 unweighted cases and have been suppressed. [1]Skilled provider includes doctor, nurse, or midwife. [2]Fully vaccinated includes BCG, measles, three doses each of DPT and polio vaccine (excluding polio vaccine given at birth) and pneumococcal. [3]An insecticide-treated net (ITN) is (1) a factory-treated net that does not require any further treatment, or (2) a net that has been soaked with insecticide within the past 6 months.

Nyanza: Key Indicators from the 2014 KDHS


Total Fertility Rate

Average number of live births per woman age 15-49




Use of Modern Methods of Family Planning

Percent of currently married women using a modern method of contraception




Antenatal Care: 4+ visits

Among women with a live birth in the 5 years before the survey, percent with 4+ ANC visits for their last live birth




Delivery in a Health Facility

Percentage of live births delivered in a health facility




Full Vaccination Rate

Percent of children age 12-23 months who have received all basic vaccinations plus the pneumococcal vaccine



ITN Access

Percentage of households with at least one insecticide-treated net (ITN) for every two persons who stayed in the household last night



The 2014 Kenya Demographic and Health Survey (2014 KDHS) was implemented by the Kenya National Bureau of Statistics from May 2014 to October 2014 in partnership with the Ministry of Health, the National AIDS Control Council (NACC), the National Council for Population and Development (NCPD), and the Kenya Medical Research Institute (KEMRI). Funding for the KDHS was provided by the Government of Kenya with support from the United States Agency for International Development (USAID), the United Nations Population Fund (UNFPA), the United Kingdom Department for International Development (DfID), the World Bank, the Danish International Development Agency (DANIDA), the United Nations Children's Fund (UNICEF), the German Development Bank (KfW), the Clinton Health Access Initiative (CHAI), the World Food Programme (WFP), and the Micronutrient Initiative (MI). ICF International provided technical assistance as well as funding to the project through The DHS Program, a USAID-funded project providing support and technical assistance in the implementation of population and health surveys in countries worldwide.

Republic of Kenya

Additional information about the 2014 KDHS may be obtained from the Kenya National Bureau of Statistics (KNBS), P.O. Box 30266-00100 GPO Nairobi, Kenya; telephone (Nairobi): 3317586/8, 3317612/22, 3317623, 3317651; fax: 3315977; e-mail: directorgeneral@knbs.or.ke, info@knbs.or.ke; website: www.knbs.or.ke.

KNBS KENYA NATIONAL BUREAU OF STATISTICS *Keeping you informed*

Information on The DHS Program may be obtained from ICF International, 530 Gaither Road, Suite 500, Rockville MD, 20850, USA; telephone: 301-407-6500; fax: 301.407.6501; e-mail: info@DHSprogram.com; website: www.DHSprogram.com.



2014 Kenya DHS
Demographic and Health Survey

Nairobi


	Nairobi	Kenya total
Fertility and Family Planning		
Total fertility rate (number of children per woman)	2.7	3.9
Current use of *any* method of family planning (% of currently married women age 15-49)	63	58
Current use of any modern method of family planning (% of currently married women age 15-49)	58	53
Maternal and Child Health		
Maternity care		
Pregnant women who received antenatal care from a skilled provider[1] (%)	98	96
Pregnant women who received 4+ ANC visits (%)	73	58
Births assisted by a skilled provider[1] (%)	89	62
Births delivered in a health facility (%)	89	61
Child vaccination		
Children 12–23 months fully vaccinated[2] (%)	60	68
Nutrition		
Children under 5 years who are stunted (moderate or severe) (%)	17	26
Children under 5 years who are wasted (moderate or severe) (%)	3	4
Children under 5 years who are underweight (%)	4	11
Children under 5 years who are overweight or obese (%)	5	4
Malaria Prevention and Treatment		
Households with at least one insecticide-treated net[3] (ITN) (%)	43	59
Households with at least one ITN for every 2 persons in the household (universal coverage) (%)	29	34
Children under age 5 who slept under an ITN the night before the survey (%)	50	54
Pregnant women who slept under an ITN the night before the survey (%)	43	51
Pregnant women who received 3+ doses of SP/Fansidar to prevent malaria during pregnancy, at least one dose was received during an ANC visit (IPTp) (%)	0.8	10
Children under age 5 with fever in the 2 weeks before the survey	19	24
Among children under age 5 with fever, percentage who took any ACT	7	23
HIV/AIDS-related Knowledge		
Women 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	84	77
Men 15-49 who know that HIV can be prevented by using condoms and limiting sex to one uninfected partner (%)	90	85
Women 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	60	53
Men 15-49 who have been tested for HIV in the past 12 months and received the results of the last test (%)	58	46

Note: Figures in parentheses are based on 25-49 unweighted cases; an asterisk indicates that data are based on fewer than 25 unweighted cases and have been suppressed. [1]Skilled provider includes doctor, nurse, or midwife. [2]Fully vaccinated includes BCG, measles, three doses each of DPT and polio vaccine (excluding polio vaccine given at birth) and pneumococcal. [3]An insecticide-treated net (ITN) is (1) a factory-treated net that does not require any further treatment, or (2) a net that has been soaked with insecticide within the past 6 months.

Central and Nairobi: Key Indicators from the 2014 KDHS


Total Fertility Rate

Average number of live births per woman age 15-49




Use of Modern Methods of Family Planning

Percent of currently married women using a modern method of contraception




Antenatal Care: 4+ visits

Among women with a live birth in the 5 years before the survey, percent with 4+ ANC visits for their last live birth




Delivery in a Health Facility

Percentage of live births delivered in a health facility




Full Vaccination Rate

Percent of children age 12-23 months who have received all basic vaccinations plus the pneumococcal vaccine



ITN Access

Percentage of households with at least one insecticide-treated net (ITN) for every two persons who stayed in the household last night



The 2014 Kenya Demographic and Health Survey (2014 KDHS) was implemented by the Kenya National Bureau of Statistics from May 2014 to October 2014 in partnership with the Ministry of Health, the National AIDS Control Council (NACC), the National Council for Population and Development (NCPD), and the Kenya Medical Research Institute (KEMRI). Funding for the KDHS was provided by the Government of Kenya with support from the United States Agency for International Development (USAID), the United Nations Population Fund (UNFPA), the United Kingdom Department for International Development (DfID), the World Bank, the Danish International Development Agency (DANIDA), the United Nations Children's Fund (UNICEF), the German Development Bank (KfW), the Clinton Health Access Initiative (CHAI), the World Food Programme (WFP), and the Micronutrient Initiative (MI). ICF International provided technical assistance as well as funding to the project through The DHS Program, a USAID-funded project providing support and technical assistance in the implementation of population and health surveys in countries worldwide.

Republic of Kenya

Additional information about the 2014 KDHS may be obtained from the Kenya National Bureau of Statistics (KNBS), P.O. Box 30266-00100 GPO Nairobi, Kenya; telephone (Nairobi): 3317586/8, 3317612/22, 3317623, 3317651; fax: 3315977; e-mail: directorgeneral@knbs.or.ke, info@knbs.or.ke; website: www.knbs.or.ke.

Information on The DHS Program may be obtained from ICF International, 530 Gaither Road, Suite 500, Rockville MD, 20850, USA; telephone: 301-407-6500; fax: 301.407.6501; e-mail: info@DHSprogram.com; website: www.DHSprogram.com.